
TUM/MPP Collider Phenomenology Seminar

Towards QED at N³LO


Yannick Ulrich

AEC, University of Bern


3 JULY 2024

- where & why do we need QED corrections?
- why do we need (partial) N^3LO ?
- what tools are needed for this?
- what to expect?
- some results from McMULE

most precise measurement of *g* – 2




most precise measurement of *g - 2*




	value	diagrams
QED 1-loop	$\alpha/2\pi = 11\,614\,097.3$	
QED 2-loop	-17 723.1	
QED 3-loop more QED	148.0 -0.5	
EW	15.3	
HVP	684.5(4.0)	
HLbL	9.2(1.7)	
total FNAL+BNL	11 659 181.0(4.3) 11 659 206.2(4.0)	[<i>g</i> – 2 white paper 20]

largest source of uncertainty & non-perturbative



this problem is bigger than *g* – 2! [CMD-3 23] [BMW 20]

largest source of uncertainty & non-perturbative



this problem is bigger than *g* – 2! [CMD-3 23] [BMW 20]

time-like in $ee \rightarrow \text{hadrons}$

$$\int ds \left(K(s) \begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$

space-like in $e\mu \rightarrow e\mu$

$$\int dt \left(K'(t) \begin{array}{c} \text{wavy line} \\ \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$

time-like in $ee \rightarrow \text{hadrons}$

$$\int ds \left(K(s) \begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$

space-like in $e\mu \rightarrow e\mu$

$$\int dt \left(K'(t) \begin{array}{c} \text{wavy line} \\ \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$

... but what actually happens ...

radiative return measurement


$$\int ds \left(K(s) \begin{array}{c} \text{wavy line} \\ \diagup \\ \diagdown \\ \text{wavy line} \\ \hline \text{wavy line} \end{array} \right)$$

loop-induced process


$$\int dt K'(t) \left(\begin{array}{c} \text{elliptical loop} \\ - \\ \text{wavy line loop} \\ - \\ \text{wavy line loop} \end{array} \right)$$

radiative corrections are vital

time-like in $ee \rightarrow \text{hadrons}$


$$\int ds \left(K(s) \begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$


space-like in $e\mu \rightarrow e\mu$


$$\int dt \left(K'(t) \begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \end{array} \right)$$


... but what actually happens ...

radiative return measurement


$$\int ds \left(K(s) \frac{\begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \\ \hline \diagup \\ \diagdown \\ \text{wavy line} \end{array}}{\begin{array}{c} \diagup \\ \diagdown \\ \text{wavy line} \end{array}} \right)$$


loop-induced process

$$\int dt K'(t) \left(\begin{array}{c} \text{elliptical loop} \\ - \\ \text{wavy line} \\ - \\ \text{square loop} \end{array} \right)$$


radiative corrections are vital

benefiting from LHC technology where possible

- soft resummation: CEEX (\rightarrow improved YFS exponentiation)
- collinear resummation: parton shower & structure functions
- $2 \rightarrow 2$ with mass dependence at NNLO
 - \Rightarrow precision: $\mathcal{O}(10^{-4})$
 - MUonE needs 10^{-5}
- $2 \rightarrow 3$ with mass dependence at NLO
 - \Rightarrow precision: $\mathcal{O}(\text{few} \times 10^{-3})$
 - radiative return needs NNLO for kinematics
- pion final states: often only very simplified models $|F_\pi(s)|^2 \times$ 
 - full hadronic model needed

just like @ the LHC ...




just like @ the LHC ...

... except fermion masses are physical \Rightarrow need massive amplitude

- they are also small \rightarrow we can drop terms $\sim \left(\frac{\alpha}{\pi}\right)^2 \log \frac{m^2}{Q^2} \times \frac{m^2}{Q^2}$
- based on SCET factorisation & method of regions [Penin 06; Mitov, Moch 06; Becher, Melnikov 07; Engel, Gnendiger, Signer, YU 18]
- process e.g. $e\mu \rightarrow e\mu$ at two-loop:

$$\mathcal{A}(m) = \mathcal{S} \times \sqrt{Z} \times \sqrt{Z} \times \mathcal{A}(0) + \mathcal{O}(m) \supset \{1/\epsilon^2, L^2\}$$

- **soft**: process-dependent $S = 1 + \text{fermion loops}$
 \rightarrow compute separately to combine with hadron loops
- **collinear**: universal Z , converts $1/\epsilon \rightarrow \log(m^2/Q^2)$



amplitude



implementation




cross section

just like @ the LHC ...

... except the real-virtual can be delicate b/c it's more exclusive

$$= \frac{1}{E_\gamma^2}$$

$$+ \mathcal{O}(E_\gamma^{-1})$$



amplitude



implementation




cross section

just like @ the LHC ...

... except the real-virtual can be delicate b/c it's more exclusive

$$\begin{aligned}
 & \text{amplitude} = \frac{1}{E_\gamma^2} \left(\text{eikonal} + \frac{1}{E_\gamma} \left\{ D \left[\text{LBK} \right] + S \left[\text{soft function} \right] \right\} + \mathcal{O}(E_\gamma^0) \right)
 \end{aligned}$$



amplitude




implementation



cross section

test next-to-soft stabilisation vs OL4 (OpenLoops quad) for $\mu e \rightarrow \mu e$ real-virtual



- same statistics, same result
 - 70 days vs 4 days
 - integrated results for different cuts
- ⇒ this is **not** an approximation but a numerical tool

NTS	OL4
-0.29268(4)	-0.29267(4)
-0.44789(6)	-0.44778(6)
-0.64662(9)	-0.64649(9)

just like @ the LHC ...

- universal soft limit $\mathcal{M}_{n+1}^{(\ell)} = \mathcal{E} \mathcal{M}_n^{(\ell)} + \mathcal{O}(E_\gamma^{-1})$
- universal pole structure $e^{\hat{\mathcal{E}}} \sum_{\ell=0}^{\infty} \mathcal{M}_n^{(\ell)} = \sum_{\ell=0}^{\infty} \mathcal{M}_n^{(\ell)f} = \text{finite}$

use this to construct an all-order subtraction scheme FKS^ℓ [Engel, Signer, YU 19]

$$\underbrace{\int d\Phi_\gamma \text{ (diagram with grey loop)}}_{\text{divergent and complicated}} = \underbrace{\int d\Phi_\gamma \left(\text{ (diagram with grey loop)} - \text{ (diagram with green loop)} \right)}_{\text{complicated but finite}} + \underbrace{\int d\Phi_\gamma \text{ (diagram with green loop)}}_{\text{divergent but easy}}$$

amplitude



implementation



cross section

Quest for precision in hadronic cross sections at low energy: Monte Carlo tools vs. experimental data

Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies

S. Actis¹⁸, A. Arbuzov⁸, G. Balossini^{1,20}, P. Beltram¹¹, C. Bigumam^{12,13}, R. Bonciani¹⁵, C.M. Carloni Calame³⁵, V. Chervenkov²⁵, M. Czakon¹⁵, H. Czyz^{16,24}, A. Denig¹⁷, S. Eidelman^{25,26}, G.V. Fedotovich^{25,26}, A. Ferruglia²¹, J. Gluza¹⁶, A. Grzelicka¹⁶, M. Guzzi³⁷, A. Hafner²⁷, F. Ignatov²⁵, S. Jadach¹⁸, F. Jegerlehner^{19,20,21}, A. Kalinowski²¹, W. Klap¹⁷, A. Korshen²⁵, J.H. Küller³⁴, E.A. Kuravly²⁵, P. Laskin²⁵, M. Mastrolia¹⁴, G. Montagna^{32,33,4}, S.E. Müller²², F. Nguyen^{4,24}, O. Nicrion³¹, D. Nomura^{4,23}, G. Pakhlov²⁴, G. Panzeri¹¹, M. Passera³, A. Penin¹⁰, F. Piccinini¹³, W. Płaczek¹⁷, T. Przedzalski¹⁷, E. Remiddi¹¹, T. Riemann⁴¹, G. Rodriguez²⁷, P. Rojo²⁷, O. Shekhovtsova¹, C.P. Shen¹⁶, A.I. Shobudanov²⁵, T. Teubner^{30,31}, G. Venanzoni^{11,24}, J.J. van der der², P. Wang¹, B.F.L. Ward¹, Z. Wan^{4,2}, M. Worek^{10,11}, C.Z. Yuan¹

¹Institut für Theoretische Physik II, RWTH Aachen Universität, 52056 Aachen, Germany

²Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

³Institut für Theoretische Physik, 12489 Berlin, Germany

⁴Dipartimento di Fisica dell'Università di Bologna, 40126 Bologna, Italy

⁵INFN, Sezione di Bologna, 40126 Bologna, Italy

⁶The Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, Reymonta 4, 30-059 Cracow, Poland

⁷Marian Smoluchowski Institute of Physics, Jagiellonian University, Reymonta 4, 30-059 Cracow, Poland

⁸Instytut Fizyki Nuklearnej Polskiej Akademii Nauk, 31-342 Kraków, Poland

⁹Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia

¹⁰Department of Physics, University of Alberta, Edmonton, AB T6G 2P1, Canada

¹¹Laboratori Nazionali del Gran Sasso, 65124 Frascati, Italy

¹²Physik Department, CERN, 1211 Geneva, Switzerland

¹³Laboratoire de Physique Subatomique et de Cosmologie, Université Joseph Fourier/CNRS/IN2P3/DPIPG, 38026 Grenoble, France

¹⁴Laboratoire de Physique Théorique, Université Paris-Sud, 91405 Orsay Cedex, France

¹⁵Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76021 Karlsruhe, Germany

¹⁶Institut für Theoretische Teilchenphysik, Universität Karlsruhe, 76123 Karlsruhe, Germany

¹⁷Institute of Physics, University of Silesia, 40007 Katowice, Poland

¹⁸National Scientific Center "Bogoliubov Institute of Theoretical Physics", 03687 Kiev, Ukraine

¹⁹Department of Physics, Princeton University, Princeton, NJ 08544, USA

²⁰Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany

²¹Institut für Physik (Theorie), Johannes Gutenberg-Universität, 55099 Mainz, Germany

²²Institute for Theoretical and Experimental Physics, Moscow, Russia

²³Scuola Normale Superiore, 56127 Pisa, Italy

²⁴NovoSibirsk State University, 630090 Novosibirsk, Russia

²⁵Laboratoire de Physique Théorique (UMR 8627), Université de Paris-Sud XI, Bâtiment 210, 91405 Orsay Cedex, France

²⁶INFN, Sezione di Padova, 35131 Padova, Italy

²⁷ILR-Ecole Polytechnique, 91128 Palaiseau, France

²⁸Dipartimento di Fisica Nucleare e Teorica, Università di Parma, 27100 Parma, Italy

²⁹INFN, Sezione di Parma, 27100 Parma, Italy

³⁰School of Physics and Astronomy, University of Southampton, Southampton SO17 1BI, UK

³¹Theory Center, KEK, Tsukuba, Ibaraki 305-0001, Japan


³²Instituto de Física Corpuscular (IFIC), Centro mixto UV/IFIC, Edificio Institutos de Investigaciones, Apartado de Correos 22085, 46071 Valencia, Spain

³³Paul Scherrer Institut, Würenlingen and Villigen, 5232 Villigen PSI, Switzerland


³⁴Department of Physics, Baylor University, Waco, TX 76799-7316, USA


³⁵Fachbereich C, Bergische Universität Wuppertal, 42097 Wuppertal, Germany


³⁶Deutsches Elektronen-Synchrotron, DESY, 15738 Zeuthen, Germany





$e^-e^+ \rightarrow e^-e^+ @ S0.7$

















- this pipeline also works for $2 \rightarrow 3$
- ... as longs as $m_f^2 \ll s_{ij}$ (massification)

for $ee \rightarrow \mu\mu\gamma$

- photon is detected, i.e. hard & large angle
- the simplest part ($ee \rightarrow \gamma\gamma^*(\rightarrow \mu\mu)$) with $m_e^2 \ll s_{ij}$
- the full with $m_e^2 \sim m_\mu^2 \ll s_{ij}$ with amplitudes from $pp \rightarrow 2j + \gamma$ (WIP)
 \Rightarrow no theoretical showstoppers, fairly doable

VVV

- for $ee \rightarrow \gamma^* (\rightarrow \mu\mu)$: HQFF known [Fael, Lange, Schönwald, Steinhauser 22]
- for $ee \rightarrow \mu\mu$: massification (known) ✗ massless (expected)

RVV



- full mass dependence unlikely (DiffExp-style is too slow for Monte Carlo)
- massless known from three-jet production
- massification...?

RRV

- OpenLoops + NTS



NTS expansion



massification






jettification

expand for $m_e^2 \sim p_e \cdot p_\gamma \ll p_e \cdot q \sim p_\gamma \cdot q$

- calculation in SCET
- two non-trivial scales: $(p_e \cdot p_\gamma)/m_e^2$ and $(p_e \cdot q)/(p_\gamma \cdot q)$
- integrals not regularised in DIMREG

$$\frac{1}{\ell \cdot \bar{n}} \rightarrow \frac{1}{(\ell \cdot \bar{n})^{1+\eta}} \quad \text{or} \quad \frac{1}{\ell \cdot \bar{n}} \rightarrow \frac{1}{\ell \cdot \bar{n} + \Delta}$$

- either complicates the integrals
- final result J finite in η or Δ




[WIP, Schalch, Engel, YU]

- the NNLO $2 \rightarrow 2$ era has arrived, also for QED
 - NNLO $2 \rightarrow 3$ possible for many things by adapting LHC results
 - (partial) N^3LO possible in the near future
 - resummation is vital to reach the target precision
- ⇒ plans for YFS-based EEX in McMULE, PS-based in other codes
(Babayaga/MESMER)



f.l.t.r.: F.Hagelstein (Mainz), A.Coutinho (IFIC), N.Schalch (Bern), L.Naterop (Zurich & PSI),
S.Kollatzsch (Zurich & PSI), A.Signer (Zurich & PSI), M.Rocco (PSI), T.Engel (Freiburg),
V.Sharkovska (Zurich & PSI), Y.Ulrich (Bern), A.Gurgone (Pavia)
not pictured: P.Banerjee (IIT Guwahati), D.Moreno (PSI), D.Radic (PSI)



McMULE
mule-tools.gitlab.io