

Symbol Alphabets from the Landau Singular Locus

Christoph Dlapa

work with Martin Helmer, Georgios Papathanasiou and Felix Tellander

2304.02629 [hep-th] - JHEP10(2023)161



Feynman Integrals

- Momentum space representation:

$$\mathcal{I} = \int \prod_{l=1}^L \frac{d^D k_l}{i\pi^{D/2}} \int_0^\infty \prod_{e \in E} \frac{1}{(-q_e^2 + m_e^2 - i\varepsilon)^{\nu_e}}, \quad D = D_0 - 2\epsilon$$

- Master integrals and canonical differential equations:

Integration-by-parts (IBP) relations

$$d\vec{f} \xleftarrow{\text{IBP}} dM(\epsilon)\vec{f}, \quad \longrightarrow \quad d\vec{g} = \epsilon \widetilde{dM}\vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)}$$

$$\vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

Feynman Integrals

- Momentum space representation:

$$\mathcal{I} = \int \prod_{l=1}^L \frac{d^D k_l}{i\pi^{D/2}} \int_0^\infty \prod_{e \in E} \frac{1}{(-q_e^2 + m_e^2 - i\varepsilon)^\nu_e}, \quad D = D_0 - 2\epsilon$$

- Master integrals and canonical differential equations:

Integration-by-parts (IBP) relations

$$d\vec{f} = dM(\epsilon)\vec{f}, \quad \longrightarrow \quad d\vec{g} = \epsilon d\widetilde{M}\vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)}$$

- Letters and alphabet:

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i$$

Goal: Find alphabet from integral representation instead of differential equations

$$\vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

Alphabet and Bootstrap

- Symbol bootstrap

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i,$$

Alphabet and Bootstrap

- Symbol bootstrap

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i,$$

- Numerical IBP for differential equations:

Integration-by-parts (IBP) relations

$$d\vec{g} = \overset{\longleftarrow}{\epsilon} d\widetilde{M} \vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)} \quad \longrightarrow \quad \vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

[Abreu, Ita, Moriello, Page,
Tschernow, Zeng, '20 /
Henn, Matijašić, Miczajka, Peraro,
Xu, Zhang, '24]

Alphabet and Bootstrap

- Symbol bootstrap

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i,$$

- Numerical IBP for differential equations:

Integration-by-parts (IBP) relations

$$d\vec{g} = \overleftarrow{\epsilon} d\widetilde{M} \vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)} \quad \longrightarrow \quad \vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

- Canonical basis e.g. from integrand analysis

[Abreu, Ita, Moriello, Page,
Tschernow, Zeng, '20 /
Henn, Matijašić, Miczajka, Peraro,
Xu, Zhang, '24]

Alphabet and Bootstrap

- Symbol bootstrap

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i,$$

- Numerical IBP for differential equations:

Integration-by-parts (IBP) relations

$$d\vec{g} = \overleftarrow{\epsilon} d\widetilde{M} \vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)} \quad \longrightarrow \quad \vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

- Canonical basis e.g. from integrand analysis
- Used to derive DEs up to ten external legs at one loop

[Abreu, Ita, Moriello, Page, Tschernow, Zeng, '20 / Henn, Matijašić, Miczajka, Peraro, Xu, Zhang, '24]

Alphabet and Bootstrap

- Symbol bootstrap

$$\widetilde{M} = \sum_i \tilde{a}_i \log W_i,$$

- Numerical IBP for differential equations:

Integration-by-parts (IBP) relations

$$d\vec{g} = \overleftarrow{\epsilon} d\widetilde{M} \vec{g}, \quad \longrightarrow \quad \vec{g} = \sum_{k=0}^{\infty} \epsilon^k \vec{g}^{(k)} \quad \longrightarrow \quad \vec{g}^{(k)} = \int d\widetilde{M} \vec{g}^{(k-1)}$$

- Canonical basis e.g. from integrand analysis
- Used to derive DEs up to ten external legs at one loop
- Finding canonical basis: INITIAL, CANONICA

[CD, Henn, Yan, '20]

[Meyer, '17]

[Abreu, Ita, Moriello, Page, Tschernow, Zeng, '20 / Henn, Matijašić, Miczajka, Peraro, Xu, Zhang, '24]

Lee-Pomeransky Representation

- Feynman representation:

$$\mathcal{I} = \Gamma(\omega) \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{\delta(1 - H(x))}{\mathcal{U}^{D/2}} \left(\frac{1}{\mathcal{F}/\mathcal{U} - i\varepsilon} \right)^\omega, \quad \omega \equiv \sum_{e \in E} \nu_e - LD/2$$

- Lee-Pomeransky:

$$\mathcal{I} = \frac{\Gamma(D/2)}{\Gamma(D/2 - \omega)} \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{1}{\mathcal{G}^{D/2}}, \quad \mathcal{G} = \mathcal{U} + \mathcal{F}$$

Lee-Pomeransky Representation

- Feynman representation:

$$\mathcal{I} = \Gamma(\omega) \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{\delta(1 - H(x))}{\mathcal{U}^{D/2}} \left(\frac{1}{\mathcal{F}/\mathcal{U} - i\varepsilon} \right)^\omega, \quad \omega \equiv \sum_{e \in E} \nu_e - LD/2$$

- Lee-Pomeransky:

$$\begin{array}{c} 1 = \int_0^\infty \delta(t - H(x)) dt \\ x_e \rightarrow tx_e \end{array}$$

$$\mathcal{I} = \frac{\Gamma(D/2)}{\Gamma(D/2 - \omega)} \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{1}{\mathcal{G}^{D/2}}, \quad \mathcal{G} = \mathcal{U} + \mathcal{F}$$

Lee-Pomeransky Representation

- Feynman representation:

$$\mathcal{I} = \Gamma(\omega) \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{\delta(1 - H(x))}{\mathcal{U}^{D/2}} \left(\frac{1}{\mathcal{F}/\mathcal{U} - i\varepsilon} \right)^\omega, \quad \omega \equiv \sum_{e \in E} \nu_e - LD/2$$

- Lee-Pomeransky:

$$\begin{array}{c} 1 = \int_0^\infty \delta(t - H(x)) dt \\ x_e \rightarrow tx_e \end{array}$$

$$\mathcal{I} = \frac{\Gamma(D/2)}{\Gamma(D/2 - \omega)} \int_0^\infty \prod_{e \in E} \left(\frac{x_e^{\nu_e} dx_e}{x_e \Gamma(\nu_e)} \right) \frac{1}{\mathcal{G}^{D/2}}, \quad \mathcal{G} = \mathcal{U} + \mathcal{F}$$

- Landau equations:

[Klaussen, '21]

$$\mathcal{G}_h = \mathcal{U}x_0 + \mathcal{F} = 0, \quad \text{and} \quad \frac{\partial \mathcal{G}_h}{\partial x_i} = 0 \quad \text{or} \quad x_i = 0 \quad \forall i = 0, \dots, |E|$$


homogenized LP-polynomial

Generic one-loop integrals

- Landau equations:

$$\mathcal{G}_h = \mathcal{U}x_0 + \mathcal{F} = 0, \quad \text{and} \quad \frac{\partial \mathcal{G}_h}{\partial x_i} = 0 \quad \text{or} \quad x_i = 0 \quad \forall i = 0, \dots, n$$

$$n = |E|$$



Generic one-loop integrals


- Landau equations:

$$\mathcal{G}_h = \mathcal{U}x_0 + \mathcal{F} = 0, \quad \text{and} \quad \frac{\partial \mathcal{G}_h}{\partial x_i} = 0 \quad \text{or} \quad x_i = 0 \quad \forall i = 0, \dots, n$$

- leading Landau singularities (full graph): $x_e \neq 0$,

- type-I singularity $x_0 = 0 \longrightarrow \mathcal{G}_h|_{x_0=0} = \mathcal{F}$

- type-II singularity $x_0 \neq 0 \longrightarrow \mathcal{G}_h|_{x_0=1} = \mathcal{G}$



Generic one-loop integrals

- Landau equations:

$$\mathcal{G}_h = \mathcal{U}x_0 + \mathcal{F} = 0, \quad \text{and} \quad \frac{\partial \mathcal{G}_h}{\partial x_i} = 0 \quad \text{or} \quad x_i = 0 \quad \forall i = 0, \dots, n$$

$$n = |E|$$

- leading Landau singularities (full graph): $x_e \neq 0, \quad \forall e \in E$


- type-I singularity $x_0 = 0 \longrightarrow \mathcal{G}_h|_{x_0=0} = \mathcal{F}$

- type-II singularity $x_0 \neq 0 \longrightarrow \mathcal{G}_h|_{x_0=1} = \mathcal{G}$

- sub-graph singularities: $x_e = 0, \quad e \in E$

- type-I singularity $x_0 = 0 \longrightarrow \mathcal{G}_h|_{\substack{x_0=0 \\ x_e=0}} = \mathcal{F}|_{x_e=0}$

- type-II singularity $x_0 \neq 0 \longrightarrow \mathcal{G}_h|_{\substack{x_0=1 \\ x_e=0}} = \mathcal{G}|_{x_e=0}$



The Landau singular locus at one loop

- Leading type-II singularity: $x_i \neq 0 \quad \forall i = 0, \dots, n$

$$\frac{\partial \mathcal{G}_h}{\partial x_0} = \dots = \frac{\partial \mathcal{G}_h}{\partial x_n} = 0 \quad \longrightarrow \quad \mathcal{G}_h = 0$$

The Landau singular locus at one loop

- Leading type-II singularity: $x_i \neq 0 \quad \forall i = 0, \dots, n$

$$\frac{\partial \mathcal{G}_h}{\partial x_0} = \dots = \frac{\partial \mathcal{G}_h}{\partial x_n} = 0 \quad \longrightarrow \quad \mathcal{G}_h \xleftarrow{\text{degree two}} 0$$

The Landau singular locus at one loop

- Leading type-II singularity: $x_i \neq 0 \quad \forall i = 0, \dots, n$

$$\frac{\partial \mathcal{G}_h}{\partial x_0} = \dots = \frac{\partial \mathcal{G}_h}{\partial x_n} = 0 \quad \longrightarrow \quad \mathcal{G}_h = 0 \quad \text{degree two}$$

depends only on kinematics

$$\rightarrow \begin{pmatrix} \frac{\partial \mathcal{G}_h}{\partial x_0} \\ \vdots \\ \frac{\partial \mathcal{G}_h}{\partial x_n} \end{pmatrix} =: \mathcal{J}(\mathcal{G}_h) \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} = \mathbf{0},$$

The Landau singular locus at one loop

- Leading type-II singularity: $x_i \neq 0 \quad \forall i = 0, \dots, n$

$$\frac{\partial \mathcal{G}_h}{\partial x_0} = \dots = \frac{\partial \mathcal{G}_h}{\partial x_n} = 0 \quad \longrightarrow \quad \mathcal{G}_h = 0 \quad \text{degree two}$$

depends only on kinematics

$$\rightarrow \begin{pmatrix} \frac{\partial \mathcal{G}_h}{\partial x_0} \\ \vdots \\ \frac{\partial \mathcal{G}_h}{\partial x_n} \end{pmatrix} =: \mathcal{J}(\mathcal{G}_h) \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} = \mathbf{0},$$

- Solution space: $\mathbf{V}\left(\frac{\partial \mathcal{G}_h}{\partial x_0}, \dots, \frac{\partial \mathcal{G}_h}{\partial x_n}\right) := \left\{x \in (\mathbb{C}^*)^n \mid \frac{\partial \mathcal{G}_h}{\partial x_0} = \dots = \frac{\partial \mathcal{G}_h}{\partial x_n} = 0\right\}$
- Space of kinematic variables for which there is a solution:

$$\overline{\left\{ s_{ij}, m_i^2 \mid \mathbf{V}\left(\frac{\partial \mathcal{G}_h}{\partial x_0}, \dots, \frac{\partial \mathcal{G}_h}{\partial x_n}\right) \neq \emptyset \right\}} \quad \iff \quad \det(\mathcal{J}(\mathcal{G}_h)) = 0$$

The modified Cayley matrix

- For the LP-polynomial of generic one-loop integrals:

$$\mathcal{J}(\mathcal{G}_h) = \mathcal{Y}$$
$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & Y_{11} & Y_{12} & \cdots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \cdots & Y_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1n} & Y_{2n} & \cdots & Y_{nn} \end{pmatrix} \leftarrow x_0 \quad \leftarrow x_e$$

$Y_{ii} = 2m_i^2, \quad Y_{ij} = m_i^2 + m_j^2 - s_{ij-1}$

Cayley matrix

$s_{ij-1} \equiv (p_i + \dots + p_{j-1})^2$

The modified Cayley matrix

- For the LP-polynomial of generic one-loop integrals:

$$\mathcal{J}(\mathcal{G}_h) = \mathcal{Y}$$
$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & Y_{11} & Y_{12} & \cdots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \cdots & Y_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1n} & Y_{2n} & \cdots & Y_{nn} \end{pmatrix} \leftarrow \begin{array}{l} x_0 \\ \text{Cayley} \\ \text{matrix} \\ x_e \end{array}$$
$$Y_{ii} = 2m_i^2, \quad Y_{ij} = m_i^2 + m_j^2 - s_{ij-1}$$
$$s_{ij-1} \equiv (p_i + \dots + p_{j-1})^2$$

- Relation to Gram determinants

• type-II singularity: $x_0 \neq 0 \rightarrow \det(\mathcal{Y}) = -2^{n-1} G(p_1, \dots, p_n)$

Gram
determinant

The modified Cayley matrix

- For the LP-polynomial of generic one-loop integrals:

$$\mathcal{J}(\mathcal{G}_h) = \mathcal{Y}$$

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & Y_{11} & Y_{12} & \cdots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \cdots & Y_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1n} & Y_{2n} & \cdots & Y_{nn} \end{pmatrix} \leftarrow \begin{matrix} x_0 \\ x_e \end{matrix}$$

$Y_{ii} = 2m_i^2, \quad Y_{ij} = m_i^2 + m_j^2 - s_{ij-1}$
 Cayley matrix
 $s_{ij-1} \equiv (p_i + \dots + p_{j-1})^2$

- Relation to Gram determinants

- type-II singularity:

$$x_0 \neq 0 \longrightarrow \det(\mathcal{Y}) = -2^{n-1} G(p_1, \dots, p_n)$$

- type-I singularity:

$$x_0 = 0 \longrightarrow \det(Y) = (-2)^n G(q_1, \dots, q_n) \Big|_{q_i^2 = m_i^2}$$

$$G(k_1, \dots, k_m) \equiv \det_{i,j}(k_i \cdot k_j)$$

Gram determinant

Cayley determinant

The principal A-determinant at one loop

- Subgraphs correspond to diagonal minors:

$$\mathcal{Y} \begin{bmatrix} i_1 & i_2 & \cdots & i_k \\ j_1 & j_2 & \cdots & j_k \end{bmatrix} \quad \text{determinant with rows/columns removed}$$

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & Y_{11} & Y_{12} & \cdots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \cdots & Y_{2n} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & Y_{1n} & Y_{2n} & \cdots & Y_{nn} \end{pmatrix} \quad \begin{array}{l} \xleftarrow{} x_0 \\ \xleftarrow{} x_e \end{array}$$

The principal A-determinant at one loop

- Subgraphs correspond to diagonal minors:

$$\mathcal{Y} \begin{bmatrix} i_1 & i_2 & \cdots & i_k \\ j_1 & j_2 & \cdots & j_k \end{bmatrix} \quad \begin{array}{l} \text{determinant with} \\ \text{rows/columns removed} \end{array}$$

- type-I singularity:

$$\mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \det(Y)$$

- type-II singularity:

$$\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} = \det(\mathcal{Y})$$

- type-I sub-singularity:

$$\mathcal{Y} \begin{bmatrix} 1 & (e+1) \\ 1 & (e+1) \end{bmatrix} = \det(Y|_{E \setminus \{e+1\}})$$

- type-II sub-singularity:

$$\mathcal{Y} \begin{bmatrix} (e+1) \\ (e+1) \end{bmatrix} = \det(\mathcal{Y}|_{E \setminus \{e+1\}})$$

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & Y_{11} & Y_{12} & \cdots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \cdots & Y_{2n} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & Y_{1n} & Y_{2n} & \cdots & Y_{nn} \end{pmatrix} \quad \begin{array}{l} \xleftarrow{x_0} \\ \xleftarrow{x_e} \end{array}$$

The principal A-determinant at one loop

- Subgraphs correspond to diagonal minors:

$$\mathcal{Y} \begin{bmatrix} i_1 & i_2 & \cdots & i_k \\ j_1 & j_2 & \cdots & j_k \end{bmatrix} \quad \begin{array}{l} \text{determinant with} \\ \text{rows/columns removed} \end{array}$$

- type-I singularity:

$$\mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \det(Y)$$

- type-II singularity:

$$\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} = \det(\mathcal{Y})$$

- type-I sub-singularity:

$$\mathcal{Y} \begin{bmatrix} 1 & (e+1) \\ 1 & (e+1) \end{bmatrix} = \det(Y|_{E \setminus \{e+1\}})$$

- type-II sub-singularity:

$$\mathcal{Y} \begin{bmatrix} (e+1) \\ (e+1) \end{bmatrix} = \det(\mathcal{Y}|_{E \setminus \{e+1\}})$$

- (reduced) principal A-determinant:

$$\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \prod_{i=1}^{n+1} \mathcal{Y} \begin{bmatrix} i \\ i \end{bmatrix} \dots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \mathcal{Y} \begin{bmatrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{bmatrix} \prod_{i=2}^{n+1} \mathcal{Y}_{ii}$$

product of Gram and Cayley determinant of the graph and all subgraphs

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & Y_{11} & Y_{12} & \dots & Y_{1n} \\ 1 & Y_{12} & Y_{22} & \dots & Y_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1n} & Y_{2n} & \dots & Y_{nn} \end{pmatrix} \quad \begin{array}{l} \xleftarrow{} x_0 \\ \xleftarrow{} x_e \end{array}$$

Example: Bubble integral

$$\widetilde{E}_A(\mathcal{G}_h) = m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2, \quad \lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2 m_1^2 - 2p^2 m_2^2 - 2m_1^2 m_2^2$$

- The factors of the principal A-determinant give all symbol letters!

Example: Bubble integral

$$\widetilde{E}_A(\mathcal{G}_h) = m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2, \quad \lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2 m_1^2 - 2p^2 m_2^2 - 2m_1^2 m_2^2$$

- The factors of the principal A-determinant give all symbol letters!
 - square-root letters?

$$\frac{-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}} \in \{W_i\}$$

Example: Bubble integral

$$\widetilde{E}_A(\mathcal{G}_h) = m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2, \quad \lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2 m_1^2 - 2p^2 m_2^2 - 2m_1^2 m_2^2$$

- The factors of the principal A-determinant give all symbol letters!

- square-root letters?

$$\frac{-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}} \in \{W_i\}$$

➤ need to re-factorize products:

$$4m_2^2 p^2$$

Example: Bubble integral

$$\widetilde{E}_A(\mathcal{G}_h) = m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2, \quad \lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2 m_1^2 - 2p^2 m_2^2 - 2m_1^2 m_2^2$$

- The factors of the principal A-determinant give all symbol letters!

- square-root letters?

$$\frac{-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}} \in \{W_i\}$$

➤ need to re-factorize products:

[Heller, Manteuffel, Schabinger, '20]

$$4m_2^2 p^2 = \left(-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)} \right) \left(-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)} \right)$$

Example: Bubble integral

$$\widetilde{E}_A(\mathcal{G}_h) = m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2, \quad \lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2 m_1^2 - 2p^2 m_2^2 - 2m_1^2 m_2^2$$

- The factors of the principal A-determinant give all symbol letters!

- square-root letters?

$$\frac{-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}} \in \{W_i\}$$

➤ need to re-factorize products:

[Heller, Manteuffel, Schabinger, '20]

$$4m_2^2 p^2 = \left(-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)} \right) \left(-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)} \right)$$

- come from Jacobi identities:

$$-\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} = \mathcal{Y} \begin{bmatrix} 2 \\ 1 \end{bmatrix}^2 - \mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad f^2 - g = (f - \sqrt{g})(f + \sqrt{g})$$

Jacobi identities

- For odd n

$$\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \mathcal{Y} \begin{bmatrix} i \\ i \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \mathcal{Y} \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} i & j \\ i & j \end{bmatrix} = \mathcal{Y} \begin{bmatrix} i \\ i \end{bmatrix} \mathcal{Y} \begin{bmatrix} j \\ j \end{bmatrix} - \mathcal{Y} \begin{bmatrix} i \\ j \end{bmatrix}^2, \quad i \geq 2$$

- For even n

$$\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \mathcal{Y} \begin{bmatrix} i \\ i \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \mathcal{Y} \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & i & j \\ 1 & i & j \end{bmatrix} = \mathcal{Y} \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & j \\ 1 & j \end{bmatrix} - \mathcal{Y} \begin{bmatrix} 1 & i \\ 1 & j \end{bmatrix}^2$$

Jacobi identities

- For odd n

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \gamma \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} i & j \\ i & j \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} j \\ j \end{bmatrix} - \gamma \begin{bmatrix} i \\ j \end{bmatrix}^2, \quad i \geq 2$$

- For even n

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \gamma \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} \gamma \begin{bmatrix} 1 & i & j \\ 1 & i & j \end{bmatrix} = \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \gamma \begin{bmatrix} 1 & j \\ 1 & j \end{bmatrix} - \gamma \begin{bmatrix} 1 & i \\ 1 & j \end{bmatrix}^2$$

case of Gram and Cayley exchanged



Jacobi identities

- For odd $n+D_0$

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \gamma \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} i & j \\ i & j \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} j \\ j \end{bmatrix} - \gamma \begin{bmatrix} i \\ j \end{bmatrix}^2, \quad i \geq 2$$

- For even $n+D_0$

$$\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} = \gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \gamma \begin{bmatrix} i \\ 1 \end{bmatrix}^2,$$

$$\gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} \gamma \begin{bmatrix} 1 & i & j \\ 1 & i & j \end{bmatrix} = \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix} \gamma \begin{bmatrix} 1 & j \\ 1 & j \end{bmatrix} - \gamma \begin{bmatrix} 1 & i \\ 1 & j \end{bmatrix}^2$$

case of Gram and Cayley exchanged



Symbol letters

- Case of one edge missing: (next-to-maximal cut)

$$W_{1,\dots,(i-1),\dots,n} = \begin{cases} \frac{\gamma \begin{bmatrix} i \\ 1 \end{bmatrix} - \sqrt{-\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix}}}{\gamma \begin{bmatrix} i \\ 1 \end{bmatrix} + \sqrt{-\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix}}}, & D_0 + n \text{ odd,} \\ \frac{\gamma \begin{bmatrix} i \\ 1 \end{bmatrix} - \sqrt{\gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix}}}{\gamma \begin{bmatrix} i \\ 1 \end{bmatrix} + \sqrt{\gamma \begin{bmatrix} i \\ i \end{bmatrix} \gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix}}}, & D_0 + n \text{ even.} \end{cases}$$

Symbol letters

- Case of two edges missing: (next-to-next-to-maximal cut)

$$W_{1,\dots,(i-1),\dots,(j-1),\dots,n} = \begin{cases} \frac{\gamma \begin{bmatrix} i \\ j \end{bmatrix} - \sqrt{-\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} i & j \\ i & j \end{bmatrix}}}{\gamma \begin{bmatrix} i \\ j \end{bmatrix} + \sqrt{-\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \gamma \begin{bmatrix} i & j \\ i & j \end{bmatrix}}}, & D_0 + n \text{ odd,} \\ \frac{\gamma \begin{bmatrix} 1 & j \\ 1 & i \end{bmatrix} - \sqrt{-\gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} \gamma \begin{bmatrix} 1 & i & j \\ 1 & i & j \end{bmatrix}}}{\gamma \begin{bmatrix} 1 & j \\ 1 & i \end{bmatrix} + \sqrt{-\gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix} \gamma \begin{bmatrix} 1 & i & j \\ 1 & i & j \end{bmatrix}}}, & D_0 + n \text{ even,} \end{cases}$$

Symbol letters

- Case of no leg missing: (maximal cut)

- no Jacobi identities
 - only one letter

$$W_{1,2,\dots,n} = \frac{\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix}}{\mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}$$

$$\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \prod_{i=1}^{n+1} \mathcal{Y} \begin{bmatrix} i \\ i \end{bmatrix} \dots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \mathcal{Y} \begin{bmatrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{bmatrix} \prod_{i=2}^{n+1} \mathcal{Y}_{ii}$$

Symbol letters

- Case of no leg missing: (maximal cut)

- no Jacobi identities
 - only one letter

$$W_{1,2,\dots,n} = \frac{\gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix}}{\gamma \begin{bmatrix} 1 \\ 1 \end{bmatrix}}$$

$$\widetilde{E}_A(\mathcal{G}_h) = \gamma \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \prod_{i=1}^{n+1} \gamma \begin{bmatrix} i \\ i \end{bmatrix} \dots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \gamma \begin{bmatrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{bmatrix} \prod_{i=2}^{n+1} \gamma_{ii}$$

- Letters not all independent

- triangle in even dimensions:

$$\log W_{(i),j,(k)} = \log W_{(i),(j),k} + \log W_{i,(j),(k)}$$

$$\frac{a - b - c - \sqrt{\lambda}}{a - b - c + \sqrt{\lambda}} = \frac{a - b + c - \sqrt{\lambda}}{a - b + c + \sqrt{\lambda}} \frac{a + b - c - \sqrt{\lambda}}{a + b - c + \sqrt{\lambda}}$$

Differential equations:

- For even $n + D_0$

$$\begin{aligned} d\mathcal{J}_{1\dots n} = & \epsilon d \log W_{1\dots n} \mathcal{J}_{1\dots n} \\ & + \epsilon \sum_{1 \leq i \leq n} (-1)^{i+\lfloor \frac{n}{2} \rfloor} d \log W_{1\dots(i)\dots n} \mathcal{J}_{1\dots \hat{i}\dots n} \\ & + \epsilon \sum_{1 \leq i < j \leq n} (-1)^{i+j+\lfloor \frac{n}{2} \rfloor} d \log W_{1\dots(i)\dots(j)\dots n} \mathcal{J}_{1\dots \hat{i}\dots \hat{j}\dots n}, \end{aligned}$$

- For odd $n + D_0$

$$\begin{aligned} d\mathcal{J}_{1\dots n} = & \epsilon d \log W_{1\dots n} \mathcal{J}_{1\dots n} \\ & + \epsilon \sum_{1 \leq i \leq n} (-1)^{i+\lfloor \frac{n+1}{2} \rfloor} d \log W_{1\dots(i)\dots n} \mathcal{J}_{1\dots \hat{i}\dots n} \\ & + \epsilon \sum_{1 \leq i < j \leq n} (-1)^{i+j+\lfloor \frac{n+1}{2} \rfloor} d \log W_{1\dots(i)\dots(j)\dots n} \mathcal{J}_{1\dots \hat{i}\dots \hat{j}\dots n}, \end{aligned}$$

Canonical master integrals

- From literature

$$\mathcal{J}_{i_1 \dots i_k} = \begin{cases} \frac{\epsilon^{\lfloor \frac{k}{2} \rfloor} \mathcal{I}_{i_1 \dots i_k}^{(k)}}{j_{i_1 \dots i_k}} & \text{for } k + D_0 \text{ even,} \\ \frac{\epsilon^{\lfloor \frac{k+1}{2} \rfloor} \mathcal{I}_{i_1 \dots i_k}^{(k+1)}}{j_{i_1 \dots i_k}} & \text{for } k + D_0 \text{ odd,} \end{cases}$$

dimension of the integral
⇒ DRR to D_0

[Abreu, Britto, Duhr, Gardi, '17 /
Chen, Ma, Yang, '22]

Canonical master integrals

- From literature

$$\mathcal{J}_{i_1 \dots i_k} = \begin{cases} \frac{\epsilon^{\lfloor \frac{k}{2} \rfloor} \mathcal{I}_{i_1 \dots i_k}^{(k)}}{j_{i_1 \dots i_k}} & \text{for } k + D_0 \text{ even,} \\ \frac{\epsilon^{\lfloor \frac{k+1}{2} \rfloor} \mathcal{I}_{i_1 \dots i_k}^{(k+1)}}{j_{i_1 \dots i_k}} & \text{for } k + D_0 \text{ odd,} \end{cases}$$

dimension of the integral
⇒ DRR to D_0

[Abreu, Britto, Duhr, Gardi, '17 /
Chen, Ma, Yang, '22]

- Leading singularities

$$j_{i_1 \dots i_k} = \begin{cases} 2^{-\frac{k}{2}+1} \left[(-1)^{\lfloor \frac{k}{2} \rfloor} \mathcal{Y} \begin{pmatrix} i_1 + 1 & i_2 + 1 & \cdots & i_k + 1 \\ i_1 + 1 & i_2 + 1 & \cdots & i_k + 1 \end{pmatrix} \right]^{-1/2}, & \text{for } k + D_0 \text{ even,} \\ 2^{-\frac{k+1}{2}+1} \left[(-1)^{\lfloor \frac{k+1}{2} \rfloor} \mathcal{Y} \begin{pmatrix} 1 & i_1 + 1 & i_2 + 1 & \cdots & i_k + 1 \\ 1 & i_1 + 1 & i_2 + 1 & \cdots & i_k + 1 \end{pmatrix} \right]^{-1/2}, & \text{for } k + D_0 \text{ odd.} \end{cases}$$

Gram determinant

Cayley determinant

Example: bubble integral $n = 2$

- Modified Cayley matrix:

$$\mathcal{G} = x_1 + x_2 + (m_1^2 + m_2^2 - p^2)x_1x_2 + m_1^2x_1^2 + m_2^2x_2^2,$$

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 2m_1^2 & m_1^2 + m_2^2 - p^2 \\ 1 & m_1^2 + m_2^2 - p^2 & 2m_2^2 \end{pmatrix}.$$

- Principal A-determinant: $\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 3 \\ 3 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix},$
 $= m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2,$

Example: bubble integral $n = 2$

- Modified Cayley matrix:

$$\mathcal{G} = x_1 + x_2 + (m_1^2 + m_2^2 - p^2)x_1x_2 + m_1^2x_1^2 + m_2^2x_2^2,$$

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 2m_1^2 & m_1^2 + m_2^2 - p^2 \\ 1 & m_1^2 + m_2^2 - p^2 & 2m_2^2 \end{pmatrix}.$$

- Principal A-determinant: $\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 3 \\ 3 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix},$
- Even D_0
 $= m_1^2 m_2^2 \lambda(p^2, m_1^2, m_2^2) p^2,$

$$W_1 = \frac{\mathcal{Y} \begin{bmatrix} 3 \\ 3 \end{bmatrix}}{\mathcal{Y} \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix}} = \frac{-1}{2m_1^2}, \quad W_2 = \frac{\mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix}}{\mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}} = \frac{-1}{2m_2^2}, \quad W_{12} = \frac{\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix}}{\mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} = \frac{2p^2}{\lambda(p^2, m_1^2, m_2^2)},$$

$$\lambda(p^2, m_1^2, m_2^2) = p^4 + m_1^4 + m_2^4 - 2p^2m_1^2 - 2p^2m_2^2 - 2m_1^2m_2^2$$

Example: bubble integral $n = 2$

- Modified Cayley matrix:

$$\mathcal{G} = x_1 + x_2 + (m_1^2 + m_2^2 - p^2)x_1x_2 + m_1^2x_1^2 + m_2^2x_2^2,$$

- Even D_0

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 2m_1^2 & m_1^2 + m_2^2 - p^2 \\ 1 & m_1^2 + m_2^2 - p^2 & 2m_2^2 \end{pmatrix}.$$

$$W_{(1)2} = \frac{\mathcal{Y} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \sqrt{\mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}}{\mathcal{Y} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \sqrt{\mathcal{Y} \begin{bmatrix} 2 \\ 2 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}} = \frac{-m_1^2 + m_2^2 + p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 + p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}},$$

$$W_{1(2)} = \frac{\mathcal{Y} \begin{bmatrix} 3 \\ 1 \end{bmatrix} - \sqrt{\mathcal{Y} \begin{bmatrix} 3 \\ 3 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}}{\mathcal{Y} \begin{bmatrix} 3 \\ 1 \end{bmatrix} + \sqrt{\mathcal{Y} \begin{bmatrix} 3 \\ 3 \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}} = \frac{-m_1^2 + m_2^2 - p^2 - \sqrt{\lambda(p^2, m_1^2, m_2^2)}}{-m_1^2 + m_2^2 - p^2 + \sqrt{\lambda(p^2, m_1^2, m_2^2)}},$$

Example: bubble integral $n = 2$

- Modified Cayley matrix:

$$\mathcal{G} = x_1 + x_2 + (m_1^2 + m_2^2 - p^2)x_1x_2 + m_1^2x_1^2 + m_2^2x_2^2,$$

- Even D_0

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 2m_1^2 & m_1^2 + m_2^2 - p^2 \\ 1 & m_1^2 + m_2^2 - p^2 & 2m_2^2 \end{pmatrix}$$

- Differential equation: $d\vec{g} = \epsilon d\widetilde{M}\vec{g},$

$$\widetilde{M} = \begin{pmatrix} w_1 & 0 & 0 \\ 0 & w_2 & 0 \\ -w_{1(2)} & w_{(1)2} & w_{12} \end{pmatrix}, \quad w = \log W$$

Example: bubble integral $n = 2$

- Modified Cayley matrix:

$$\mathcal{G} = x_1 + x_2 + (m_1^2 + m_2^2 - p^2)x_1x_2 + m_1^2x_1^2 + m_2^2x_2^2,$$

- Odd D_0

$$\mathcal{Y} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 2m_1^2 & m_1^2 + m_2^2 - p^2 \\ 1 & m_1^2 + m_2^2 - p^2 & 2m_2^2 \end{pmatrix}.$$

$$W_{(1)2} = \frac{\mathcal{Y} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \sqrt{-\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}}}{\mathcal{Y} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \sqrt{-\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}}} = \frac{-m_1^2 + m_2^2 + p^2 - \sqrt{4p^2m_2^2}}{-m_1^2 + m_2^2 + p^2 + \sqrt{4p^2m_2^2}},$$

$$W_{1(2)} = \frac{\mathcal{Y} \begin{bmatrix} 3 \\ 1 \end{bmatrix} - \sqrt{-\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix}}}{\mathcal{Y} \begin{bmatrix} 3 \\ 1 \end{bmatrix} + \sqrt{-\mathcal{Y} \begin{bmatrix} \cdot \\ \cdot \end{bmatrix} \mathcal{Y} \begin{bmatrix} 1 & 3 \\ 1 & 3 \end{bmatrix}}} = \frac{-m_1^2 + m_2^2 - p^2 - \sqrt{4p^2m_1^2}}{-m_1^2 + m_2^2 - p^2 + \sqrt{4p^2m_1^2}}$$

Comparison with the literature

- Symbol alphabet
- Differential equations

1) Diagrammatic coaction:

[Abreu, Britto, Duhr, Gardi, '17]



only next-to-next-to
maximal cut needed

also: [Caron-Huot,
Pokraka, '21]

2) Baikov representation:

[Chen, Ma, Yang, '22]



explicit match

[Jiang, Yang, '23]



modified Cayley matrix

[Jiang, Liu, Xu, Yang, '24]



higher loops

Limits to non-generic cases

$$\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \left[\frac{\cdot}{\cdot} \right] \prod_{i=1}^{n+1} \mathcal{Y} \left[\begin{matrix} i \\ i \end{matrix} \right] \dots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \mathcal{Y} \left[\begin{matrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{matrix} \right] \prod_{i=2}^{n+1} \mathcal{Y}_{ii}$$

- Consider limits, e.g. $m_i^2, s_{ij}^2 \rightarrow 0$
 - leading term in Taylor expansion

Limits to non-generic cases

$$\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \left[\begin{matrix} \cdot \\ \cdot \\ \cdot \end{matrix} \right] \prod_{i=1}^{n+1} \mathcal{Y} \left[\begin{matrix} i \\ i \end{matrix} \right] \cdots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \mathcal{Y} \left[\begin{matrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{matrix} \right] \prod_{i=2}^{n+1} \mathcal{Y}_{ii}$$

- Consider limits, e.g. $m_i^2, s_{ij}^2 \rightarrow 0$
 - leading term in Taylor expansion
 - multivariate limit for individual factors is not unique,
however, limit of principle A-determinant is!

Limits to non-generic cases

$$\widetilde{E}_A(\mathcal{G}_h) = \mathcal{Y} \left[\frac{\cdot}{\cdot} \right] \prod_{i=1}^{n+1} \mathcal{Y} \left[\begin{matrix} i \\ i \end{matrix} \right] \dots \prod_{i_{n-1} > \dots > i_1 = 1}^{n+1} \mathcal{Y} \left[\begin{matrix} i_1 \dots i_{n-1} \\ i_1 \dots i_{n-1} \end{matrix} \right] \prod_{i=2}^{n+1} \mathcal{Y}_{ii}$$

- Consider limits, e.g. $m_i^2, s_{ij}^2 \rightarrow 0$
 - leading term in Taylor expansion
 - multivariate limit for individual factors is not unique, however, limit of principle A-determinant is!
- Limits match direct computation
 - expect also to work for symbol alphabet

[Abreu, Britto, Duhr, Gardi, '17]

[Chen, Ma, Yang, '22]

Higher loops

- Principal A-determinant:

$$\overline{\left\{ s_{ij}, m_i^2 \mid \mathbf{V} \left(x_0 \frac{\partial \mathcal{G}_h}{\partial x_0}, \dots, x_n \frac{\partial \mathcal{G}_h}{\partial x_n} \right) \neq \emptyset \right\}} \iff \widetilde{E}_A(\mathcal{G}_h) = 0$$

Higher loops

- Principal A-determinant:

$$\overline{\left\{ s_{ij}, m_i^2 \mid \mathbf{V} \left(x_0 \frac{\partial \mathcal{G}_h}{\partial x_0}, \dots, x_n \frac{\partial \mathcal{G}_h}{\partial x_n} \right) \neq \emptyset \right\}} \iff \widetilde{E}_A(\mathcal{G}_h) = 0$$

- Prime factorization:

$$\widetilde{E}_A(\mathcal{G}_h) = \prod_{\Gamma \subseteq \text{Newt}(\mathcal{G}_h)} \Delta_{A \cap \Gamma}(\mathcal{G}_h|_{\Gamma}),$$

faces

A-discriminants

restriction of \mathcal{G}_h on Γ

$$\mathcal{G}_h = \sum_{i=0}^r c_i x^{\alpha_i}$$

$A = \text{Supp}(\mathcal{G}_h)$
 $\text{Newt}(\mathcal{G}_h) = \text{conv}(A)$

Higher loops

- Principal A-determinant:

$$\overline{\left\{ s_{ij}, m_i^2 \mid \mathbf{V} \left(x_0 \frac{\partial \mathcal{G}_h}{\partial x_0}, \dots, x_n \frac{\partial \mathcal{G}_h}{\partial x_n} \right) \neq \emptyset \right\}}$$

PLD: [Fevola, Mizera, Telen, '24]

Baikov: [Caron-Huot, Correia, Giroux, '24]

- Prime factorization:

$$\widetilde{E}_A(\mathcal{G}_h) = \prod_{\Gamma \subseteq \text{Newt}(\mathcal{G}_h)} \Delta_{A \cap \Gamma}(\mathcal{G}_h|_\Gamma),$$

faces



$$\mathcal{G}_h = \sum_{i=0}^r c_i x^{\alpha_i}$$

$$A = \text{Supp}(\mathcal{G}_h)$$

$$\text{Newt}(\mathcal{G}_h) = \text{conv}(A)$$

- singularities: type-I

$$\Delta_{A \cap \text{Newt}(\mathcal{F})}(\mathcal{F}),$$

- type-II

$$\Delta_A(\mathcal{G}),$$

- mixed

$$\Delta_{A \cap \Gamma}(\mathcal{G}|_\Gamma),$$


vertices on both $\text{Supp}(\mathcal{U})$ and $\text{Supp}(\mathcal{F})$

Slashed box example

- One-mass configuration

$$p_1^2 \neq 0, \quad m_i^2 = p_2^2 = p_3^2 = p_4^2 = 0$$

$$\widetilde{E}_A(\mathcal{G}_h) = (p_1^2 - t)(p_1^2 - s)(p_1^2 - s - t)(s + t)st p_1^2$$



Slashed box example

- One-mass configuration


$$p_1^2 \neq 0, \quad m_i^2 = p_2^2 = p_3^2 = p_4^2 = 0$$

$$\widetilde{E}_A(\mathcal{G}_h) = (p_1^2 - t)(p_1^2 - s)(p_1^2 - s - t)(s + t)st p_1^2$$

- two-dimensional harmonic polylogarithms:

$$z_1 = \frac{s}{p_1^2}, z_2 = \frac{t}{p_1^2}, z_3 = 1 - z_1 - z_2$$

$$\widetilde{E}_A(\mathcal{G}_h) \propto (1 - z_2)(1 - z_1)z_3(1 - z_3)z_1z_2$$



Summary

- Construction of symbol alphabet from the principal A-determinant
 - rational letters
 - square-root letters through re-factorization
- One loop
 - re-factorization through Jacobi identities
 - verification through canonical DEs (up to ten legs)
- Unique limits
- Higher loops