

# Introduction to the Standard Model

IMPRS Elementary Particle Physics

MPP, October 28 - 31, 2013



W. HOLLIK



MAX-PLANCK-GESELLSCHAFT

MAX-PLANCK-INSTITUT FÜR PHYSIK, MÜNCHEN

# Outline

1. Elements of Quantum Field Theory
2. Gauge Theories
3. QCD
4. Higgs mechanism
5. Electroweak interaction and Standard Model
6. Phenomenology of  $W$  and  $Z$  bosons, precision tests
7. Higgs bosons

# Notations and Conventions

$$\mu, \nu, .. = 0, 1, 2, 3; \quad k, l, .. = 1, 2, 3$$

$$x = (x^\mu) = (x^0, \vec{x}), \quad x^0 = t \quad (\hbar = c = 1)$$

$$p = (p^\mu) = (p^0, \vec{p}), \quad p^0 = E = \sqrt{\vec{p}^2 + m^2}$$

$$a_\mu = g_{\mu\nu} a^\nu, \quad (g_{\mu\nu}) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

$$a^2 = a_\mu a^\mu, \quad a \cdot b = a_\mu b^\mu = a^0 b^0 - \vec{a} \cdot \vec{b}$$

$$\partial_\mu = \frac{\partial}{\partial x^\mu} = g_{\mu\nu} \partial^\nu, \quad \partial^\nu = \frac{\partial}{\partial x_\nu} \quad [ \partial^0 = \partial_0, \quad \partial^k = -\partial_k ]$$

$$\square = \partial_\mu \partial^\mu = \frac{\partial^2}{\partial t^2} - \Delta$$

## **2. Gauge theories**

# Constructing QED – main steps

- start with  $\mathcal{L}_0 = \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi$  for free fermion field  $\psi$   
symmetric under global gauge transformations  
 $\psi' = e^{i\alpha} \psi$ ,  $\alpha$  real

- perform minimal substitution  $\partial_\mu \rightarrow D_\mu = \partial_\mu - ieA_\mu$   
 $\Rightarrow$  invariance under local gauge transformations  
 $\psi' = e^{i\alpha(x)} \psi$ ,  $A'_\mu = A_\mu + \frac{1}{e} \partial_\mu \alpha(x)$

- involves additional vector field  $A_\mu$
- induces interaction between  $A_\mu$  and  $\psi$

$$e (\bar{\psi} \gamma^\mu \psi) A_\mu \equiv e j^\mu A_\mu$$

- make  $A_\mu$  a dynamical field by adding

$$\mathcal{L}_A = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

# Non-Abelian gauge theories

Generalization: “phase” transformations that do not commute

$$\psi \rightarrow \psi' = U\psi \quad \text{with} \quad U_1 U_2 \neq U_2 U_1$$

requires **matrices**, *i.e.*  $\psi$  is a multiplet

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \vdots \\ \psi_n \end{pmatrix}, \quad U = n \times n \text{-matrix}$$

each  $\psi_k = \psi_k(x)$  is a Dirac spinor

## (i) global symmetry

starting point:  $\mathcal{L}_0 = \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi$

where  $\bar{\psi} = (\bar{\psi}_1, \dots, \bar{\psi}_n)$

consider unitary matrices:  $U^\dagger = U^{-1}$

$$\psi' = U\psi \quad \Rightarrow \quad \bar{\psi}' = \bar{\psi} U^\dagger = \bar{\psi} U^{-1}$$

$$\Rightarrow \quad \bar{\psi}' \psi' = \bar{\psi} \psi, \quad \bar{\psi}' \gamma^\mu \partial_\mu \psi' = \bar{\psi} \gamma^\mu \partial_\mu \psi$$

*if  $U$  does not depend on  $x$*

$\Rightarrow \mathcal{L}_0$  is invariant under  $\psi \rightarrow U\psi$

**$U$ : global gauge transformation**

similar for

scalar fields:

$$\phi \rightarrow \phi' = U\phi, \quad \phi = \begin{pmatrix} \phi_1 \\ \phi_2 \\ \vdots \\ \phi_n \end{pmatrix}$$

each  $\psi_k = \phi_k(x)$  is a scalar field,  $\phi^\dagger = (\phi_1^\dagger, \dots, \phi_n^\dagger)$

terms  $\phi^\dagger\phi$ ,  $(\partial_\mu\phi)^\dagger(\partial^\mu\phi)$  are invariant

$\Rightarrow \mathcal{L}_0 = (\partial_\mu\phi)^\dagger(\partial^\mu\phi) - m^2\phi^\dagger\phi$  is invariant



relevant in physics:

*the special unitary  $n \times n$ -matrices with  $\det=1$*

group  $SU(n)$

examples:

$$SU(2) : \quad \psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad \text{e.g.} \quad \psi = \begin{pmatrix} \psi_\nu \\ \psi_e \end{pmatrix} \quad \textit{isospin}$$

$$SU(3) : \quad \psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} \quad \text{e.g.} \quad \psi = \begin{pmatrix} \psi_r \\ \psi_g \\ \psi_b \end{pmatrix} \quad \textit{colour}$$

$SU(n)$  matrices  $U$  can be written as exponentials

$$U(\theta_1, \dots, \theta_N) = e^{i\theta_a T_a} \quad \text{sum over } a = 1, \dots, N$$

$\theta_1, \dots, \theta_N$  : real parameters

$T_1, \dots, T_N$  :  $n \times n$ -matrices, generators,  $T_a^\dagger = T_a$

infinitesimal  $\theta$  :  $U = \mathbf{1} + i\theta_a T_a \quad (+O(\theta^2))$

N-dimensional Lie Group

det=1 and unitarity  $\Rightarrow$   $N = n^2 - 1$

$n = 2$  :  $N = 3$ ,       $n = 3$  :  $N = 8$

commutators  $[T_a, T_b] \neq 0$  non-Abelian

$$\boxed{[T_a, T_b] = i f_{abc} T_c}$$

*Lie Algebra*

$f_{abc}$  : real numbers, **structure constants**

$f_{abc} = -f_{bac} = \dots$  antisymmetric

commutators  $[T_a, T_b] \neq 0$  non-Abelian

$$[T_a, T_b] = i f_{abc} T_c$$

*Lie Algebra*

$f_{abc}$  : real numbers, **structure constants**

$f_{abc} = -f_{bac} = \dots$  antisymmetric

$SU(2)$   $f_{abc} = \epsilon_{abc}$  (like angular momentum)

$T_a = \frac{1}{2} \sigma_a$ ,  $\sigma_a$  : Pauli matrices ( $a=1,2,3$ )

commutators  $[T_a, T_b] \neq 0$  non-Abelian

$$[T_a, T_b] = f_{abc} T_c$$

*Lie Algebra*

$f_{abc}$  : real numbers, **structure constants**

$f_{abc} = -f_{bac} = \dots$  antisymmetric

$SU(2)$   $f_{abc} = \epsilon_{abc}$  (like angular momentum)

$T_a = \frac{1}{2} \sigma_a$ ,  $\sigma_a$  : Pauli matrices ( $a=1,2,3$ )

$SU(3)$   $T_a = \frac{1}{2} \lambda_a$ ,  $\lambda_a$  : Gell-Mann matrices ( $a=1, \dots, 8$ )

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \dots$$

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix},$$

$$\lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}$$

## (ii) local symmetry

now:  $\theta_a = \theta_a(x)$  for  $a = 1, \dots, N$

covariant derivative  $\partial_\mu \rightarrow D_\mu = \partial_\mu - ig \mathbf{W}_\mu$

vector field  $\mathbf{W}_\mu$  is  $n \times n$  matrix:  $\mathbf{W}_\mu(x) = T_a W_\mu^a(x)$

induces interaction term  $\mathcal{L}_0 \rightarrow \mathcal{L}_0 + \mathcal{L}_{\text{int}}$

with  $\mathcal{L}_{\text{int}} = g \bar{\Psi} \gamma^\mu \mathbf{W}_\mu \Psi = g (\bar{\Psi} \gamma^\mu T_a \Psi) W_\mu^a \equiv j_a^\mu W_\mu^a$

For a multiplet of scalar fields  $\Phi$ :

$$\mathcal{L}_0 = (\partial_\mu \Phi)^\dagger (\partial^\mu \Phi) \quad \rightarrow \quad \mathcal{L} = (D_\mu \Phi)^\dagger (D^\mu \Phi)$$

$\mathcal{L}$  is invariant under local gauge transformations

$$\Psi \rightarrow \Psi' = U \Psi ,$$

$$\mathbf{W}_\mu \rightarrow \mathbf{W}'_\mu = U \mathbf{W}_\mu U^{-1} - \frac{i}{g} (\partial_\mu U) U^{-1}$$

for infinitesimal transformations:

$$W_\mu^a \rightarrow W'^a_\mu = W_\mu^a + \frac{1}{g} \partial_\mu \theta^a + f_{abc} W_\mu^b \theta^c$$

crucial property of covariant derivative

$$\boxed{D'_\mu U = U D_\mu}$$



### (iii) dynamics of $W_\mu^a$ fields

need: additional term  $\mathcal{L}_W \Rightarrow$  e.o.m., propagators

naive:  $\sum_a (\partial_\mu W_\nu^a - \partial_\nu W_\mu^a)^2$  *not gauge invariant*

instead: 
$$\begin{aligned} \mathbf{F}_{\mu\nu} &= D_\mu \mathbf{W}_\nu - D_\nu \mathbf{W}_\mu \equiv F_{\mu\nu}^a T_a \\ &= \partial_\mu \mathbf{W}_\nu^a - \partial_\nu \mathbf{W}_\mu^a - ig [\mathbf{W}_\mu, \mathbf{W}_\nu] \\ &= \frac{i}{g} [D_\mu, D_\nu] \end{aligned}$$

gauge transformation: 
$$\begin{aligned} \mathbf{W}_\mu &\rightarrow \mathbf{W}'_\mu, & D_\mu &\rightarrow D'_\mu \\ \Rightarrow \mathbf{F}_{\mu\nu} &\rightarrow \mathbf{F}'_{\mu\nu} = U \mathbf{F}_{\mu\nu} U^{-1} \end{aligned}$$

$$\Rightarrow \text{Tr}(\mathbf{F}'_{\mu\nu} \mathbf{F}'^{\mu\nu}) = \text{Tr}(U \mathbf{F}_{\mu\nu} U^{-1} U \mathbf{F}^{\mu\nu} U^{-1}) = \text{Tr}(\mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu})$$

*gauge invariant*

Lagrangian:

$$\mathcal{L}_W = -\frac{1}{2} \text{Tr} (\mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}) = -\frac{1}{4} \sum_a F_{\mu\nu}^a F^{a,\mu\nu}$$

components of  $\mathbf{F}_{\mu\nu}$  [using normalization  $\text{Tr}(T_a T_b) = \frac{1}{2} \delta_{ab}$  ]

$$F_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g f_{abc} W_\mu^b W_\nu^c$$

$$F_{\mu\nu}^a = \text{Abelian} + \text{non-Abelian}$$

$$\mathcal{L}_W = \underbrace{\text{quadratic}}_{\text{free part}} + \underbrace{\text{cubic} + \text{quartic}}_{\text{tri- and quadri-linear interactions}}$$

$$\begin{aligned}
\mathcal{L}_W &= -\frac{1}{4} (\partial_\mu W_\nu^a - \partial_\nu W_\mu^a)^2 \quad \Rightarrow \text{propagator} \\
&= -\frac{1}{2} g f_{abc} (\partial_\mu W_\nu^a - \partial_\nu W_\mu^a) W^{b,\mu} W^{c,\nu} \\
&= -\frac{1}{4} g^2 f_{abc} f_{ade} W_\mu^b W_\nu^c W^{d,\mu} W^{e,\nu}
\end{aligned}$$

new type of couplings:

- self-couplings of vector fields (gauge couplings)
- universal coupling constant  $g$  for fermions and vector fields

### 3. Quantum Chromodynamics (QCD)

each quark field  $q = u, d, \dots$  appears in 3 colours

$$\psi = \begin{pmatrix} q \\ q \\ q \end{pmatrix}, \quad \bar{\psi} = (\bar{q}, \bar{q}, \bar{q})$$

$$T_a = \frac{1}{2} \lambda_a \quad (a = 1, \dots, 8) \quad \text{generators, group } SU(3)$$

$$W_\mu^a \equiv G_\mu^a \quad \text{8 gluon fields}$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f_{abc} G_\mu^b G_\nu^c \quad \text{field strength}$$

$$D_\mu = \partial_\mu - ig_s T_a G_\mu^a \quad \text{covariant derivative}$$

$$g_s \quad \text{coupling constant of strong interaction,} \quad \alpha_s = \frac{g_s^2}{4\pi}$$

## QCD Lagrangian

(for one flavor of quarks)

$$\mathcal{L}_{\text{QCD}} = \mathcal{L}_{\text{F}} + \mathcal{L}_{\text{G}}, \quad \mathcal{L}_{\text{F}} = \bar{\psi} (i\gamma^\mu D_\mu - m) \psi,$$

$$\mathcal{L}_{\text{QCD}} = \bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi$$

$$+ g_s \bar{\psi} \gamma^\mu \frac{\lambda_a}{2} \psi G_\mu^a$$


$$- \frac{1}{4} G_{\mu\nu}^a G^{a,\mu\nu} - \frac{1}{2\xi} (\partial_\mu G^{a,\mu})^2$$


## QCD Lagrangian


(for one flavor of quarks)


$$\begin{aligned}\mathcal{L}_{\text{QCD}} &= \mathcal{L}_{\text{F}} + \mathcal{L}_{\text{G}} = \\ &\bar{\psi} (i\gamma^\mu \partial_\mu - m) \psi \\ &+ g_s \bar{\psi} \gamma^\mu \frac{\lambda_a}{2} \psi G_\mu^a \\ &- \frac{1}{4} G_{\mu\nu}^a G^{a, \mu\nu} - \frac{1}{2\xi} (\partial_\mu G^{a, \mu})^2\end{aligned}$$

*reality:* sum over quark flavors  $f$ , with  $\psi \rightarrow \psi_f$ ,  $m \rightarrow m_f$

- Quark-Gluon-Vertex:   $ig_S(T_a)_{kl}\gamma^\mu$

- Quark-Propagator:   $i\frac{\not{q}+m}{q^2-m^2+i\epsilon} \equiv \frac{i}{\not{q}-m+i\epsilon}$

- Gluon-Propagator:   $\frac{-ig_{\mu\nu}}{q^2+i\epsilon}$

- Triple-Gluon-Vertex: 

- Quartic-Gluon-Vertex: 

## **4. Higgs mechanism**



problem: weak interaction, gauge bosons are massive

mass terms  $\sim M^2 W_\mu^a W^{a,\mu}$  spoil local gauge invariance

- bad high energy behaviour of amplitudes and cross sections, conflict with unitarity

reason: longitudinal polarization  $\epsilon^\mu \simeq \frac{k^\mu}{M} \sim k^\mu$

- bad divergence of higher orders with loop diagrams

reason: propagators contain  $-g_{\mu\nu} + \frac{k_\mu k_\nu}{M^2}$

$\Rightarrow$  additional powers of momenta in loop integration


$\Rightarrow$  spoil renormalizability

**renormalizable theories**: divergences can be removed by a finite number of counter terms


**gauge invariant theories**: counter terms for parameters (and fields)

## W and Z are massive

- W, Z have longitudinal polarization states  
polarization vectors of W (Z)  $\epsilon_L \sim k/M_W$   
for large momentum k



bad high energy behaviour of WW scattering




bad divergence of loop integrals

way out:

new scalar with appropriate couplings to W,Z

● restoration of unitarity



● restoration UV finiteness  $\Rightarrow$  renormalizability





consistent way: Higgs mechanism  
= scalar with gauge invariant interactions  
and non-invariant ground state

# Spontaneous symmetry breaking (SSB)

physical system: has a symmetry

ground state: not symmetric




[A. Pich]

example: complex scalar field  $\phi \neq \phi^\dagger$

Lagrangian with interaction  $V$  (potential), minimum at  $\phi_0 = v$

$$\mathcal{L} = |\partial_\mu \phi|^2 - V(\phi)$$



$V = V(|\phi|)$  :  $\mathcal{L}$  symmetric under  $\phi \rightarrow e^{i\alpha} \phi$ ,  $U(1)$

$v \neq 0$  :  $\phi'_0 = e^{i\alpha} v \neq \phi_0$  not symmetric

$V = V(|\phi'_0|) = V(|\phi_0|)$  : vacuum is degenerate

write  $\phi(x) = \eta(x)e^{i\theta(x)}$ ,  $\eta$  and  $\theta$  real

$V(|\phi|) = V(\eta)$ , minimum at  $\eta = v$ :  $V'(v) = 0$ ,  $V''(v) > 0$

expand around minimum:  $\eta(x) = v + \frac{1}{\sqrt{2}} H(x)$

$$V(\eta) = V(v) + \frac{1}{2}V''(v) \cdot \frac{1}{2}H^2 + \dots$$

$$\mathcal{L} = \frac{1}{2}|\partial_\mu H|^2 - \underbrace{\frac{1}{2}V''(v) \cdot \frac{1}{2}H^2}_{m_H^2 > 0 \text{ mass of } H} + v^2|\partial_\mu \theta|^2 + \dots$$

- $H$  field is massive
- $\theta$  field is massless, no  $\theta^2$  term: **Goldstone field**
- special case of **Goldstone theorem**:

for each broken generator  $T_a$  with  $T_a \phi_0 \neq 0$

there is a massless Goldstone field  $\theta(x)$

## SSB in gauge theories

$$\mathcal{L} = (D_\mu \phi)^\dagger (D^\mu \phi) - V(|\phi|) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad D_\mu = \partial_\mu - ieA_\mu$$

invariant under local U(1) transformations:

$$\phi'(x) = e^{i\alpha(x)} \phi(x) = e^{i\alpha(x)} e^{i\theta(x)} \eta(x)$$

$$A'_\mu(x) = A_\mu(x) + \frac{1}{e} \partial_\mu \alpha(x)$$

choose  $\alpha(x) = -\theta(x)$ :  $\phi'(x) = \eta(x)$

$$\mathcal{L} = |(\partial_\mu - ieA'_\mu)\eta|^2 - V(\eta) - \frac{1}{4} F'_{\mu\nu} F'^{\mu\nu}$$

massless  $\theta$  field removed (unphysical)

$$\begin{aligned}
\mathcal{L} &= |(\partial_\mu - ieA'_\mu)(v + \frac{1}{\sqrt{2}}H)|^2 - \frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} - V \\
&= \underbrace{-\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + v^2 e^2 A'_\mu A'^\mu}_{\text{massive A-field, } m_A \sim ev} + \underbrace{\frac{1}{2}[(\partial_\mu H)^2 - m_H^2 H^2]}_{\text{neutral scalar, } m_H \neq 0} + \dots
\end{aligned}$$

in this special gauge: no Goldstone field unitary gauge

$A_\mu$ -field propagator:  $\frac{i}{k^2 - m_A^2 + i\epsilon} \underbrace{\left(-g_{\mu\nu} + \frac{k_\mu k_\nu}{m_A^2}\right)}_{\text{polarization sum of 3 pol. states}}$

massive vector field without spoiling gauge symmetry of  $\mathcal{L}$



## two different gauges

properties	$\phi$ field	$A_\mu$ field
symmetry manifest	$H, \theta$	2 polarizations (transverse)
physics manifest	$H$	3 polarizations (2 transverse + 1 longitudinal)

$\theta \rightarrow$  *longitudinal polarization of  $A_\mu$*

## **5. Electroweak Standard Model**

## preliminaries

Dirac matrices:  $\gamma^\mu$  ( $\mu = 0, 1, 2, 3$ ),  $\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2 g^{\mu\nu}$

$\bar{\Gamma} = \gamma^0 (\Gamma)^\dagger \gamma^0$ ,  $\Gamma$  any Dirac matrix oder product of matrices

further Dirac matrix:  $\gamma_5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$

$\gamma_5 \gamma^\mu + \gamma^\mu \gamma_5 = 0$ ,  $\bar{\gamma}_5 = -\gamma_5$ ,  $\gamma_5^2 = \mathbf{1}$

chiral fermions:

$\psi^L = \frac{1-\gamma_5}{2} \psi$  left-handed spinor, L-chiral spinor

$\psi^R = \frac{1+\gamma_5}{2} \psi$  right-handed spinor, R-chiral spinor

projectors on right/left chirality:  $\omega_\pm = \frac{1\pm\gamma_5}{2}$ ,  $(\omega_\pm)^2 = \omega_\pm$

## chiral currents:

$$\overline{\psi^L} \gamma^\mu \psi^L = \overline{\psi} \gamma^\mu \frac{1-\gamma_5}{2} \psi \equiv J_L^\mu \quad \text{left-handed current}$$

$$\overline{\psi^R} \gamma^\mu \psi^R = \overline{\psi} \gamma^\mu \frac{1+\gamma_5}{2} \psi \equiv J_R^\mu \quad \text{right-handed current}$$

$$J_V^\mu = \overline{\psi} \gamma^\mu \psi = J_L^\mu + J_R^\mu \quad \text{vector current}$$

$$J_A^\mu = \overline{\psi} \gamma^\mu \gamma_5 \psi = -J_L^\mu + J_R^\mu \quad \text{axialvector current}$$

## mass term:

$$m \overline{\psi} \psi = m (\overline{\psi^L} \psi^R + \overline{\psi^R} \psi^L)$$

connects  $L$  and  $R$  !

symmetry group:  $SU(2)_I \times U(1)_Y$

$SU(2)_I$  : weak isospin, generators  $T_I^a = \frac{1}{2} \sigma_a$  for  $L$ ,  $= 0$  for  $R$

$U(1)_Y$  : weak hypercharge, generator  $Y$

$$T_I^3 + Y/2 = Q$$

fermion content (ignoring possible right-handed neutrinos)

				$T_I^3$	$Y$	
leptons:	$\Psi_L^L =$	$\begin{pmatrix} \nu_e^L \\ e^L \end{pmatrix}$	$\begin{pmatrix} \nu_\mu^L \\ \mu^L \end{pmatrix}$	$\begin{pmatrix} \nu_\tau^L \\ \tau^L \end{pmatrix}$	$+\frac{1}{2}$	$-1$
	$\psi_l^R =$	$e^R$	$\mu^R$	$\tau^R$	$0$	$-2$
quarks:	$\Psi_Q^L =$	$\begin{pmatrix} u^L \\ d^L \end{pmatrix}$	$\begin{pmatrix} c^L \\ s^L \end{pmatrix}$	$\begin{pmatrix} t^L \\ b^L \end{pmatrix}$	$+\frac{1}{2}$	$+\frac{1}{3}$
	$\psi_u^R =$	$u^R$	$c^R$	$t^R$	$0$	$+\frac{4}{3}$
	$\psi_d^R =$	$d^R$	$s^R$	$b^R$	$0$	$-\frac{2}{3}$

## gauge boson content

$SU(2)_I$  : generators  $T_I^1, T_I^2, T_I^3$

gauge fields  $W_\mu^1, W_\mu^2, W_\mu^3$

also:  $W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \mp i W_\mu^2), W_\mu^3$

$U(1)_Y$  : generator  $Y$

gauge field  $B_\mu$

$$[T_I^a, T_I^b] = i\epsilon_{abc} T_I^c, \quad [T_I^a, Y] = 0$$

## gauge boson content

$SU(2)_I$  : generators  $T_I^1, T_I^2, T_I^3$

gauge fields  $W_\mu^1, W_\mu^2, W_\mu^3$

also:  $W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \mp i W_\mu^2), W_\mu^3$

$U(1)_Y$  : generator  $Y$

gauge field  $B_\mu$

notation:  $\not{\partial} = \gamma^\mu \partial_\mu, \not{a} = \gamma^\mu a_\mu$

## Free Lagrangian of (still massless) fermions:

$$\mathcal{L}_{0,\text{ferm}} = i\overline{\psi}_f \not{\partial} \psi_f = i\overline{\Psi}_L^L \not{\partial} \Psi_L^L + i\overline{\Psi}_Q^L \not{\partial} \Psi_Q^L + i\overline{\psi}_l^R \not{\partial} \psi_l^R + i\overline{\psi}_u^R \not{\partial} \psi_u^R + i\overline{\psi}_d^R \not{\partial} \psi_d^R$$

## Minimal substitution:

$$\partial_\mu \rightarrow D_\mu = \partial_\mu - ig_2 T_I^a W_\mu^a + ig_1 \frac{1}{2} Y B_\mu = D_\mu^L \omega_- + D_\mu^R \omega_+,$$

$$D_\mu^L = \partial_\mu - \frac{ig_2}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ \\ W_\mu^- & 0 \end{pmatrix} - \frac{i}{2} \begin{pmatrix} g_2 W_\mu^3 - g_1 Y^L B_\mu & 0 \\ 0 & -g_2 W_\mu^3 - g_1 Y^L B_\mu \end{pmatrix},$$

$$D_\mu^R = \partial_\mu + ig_1 \frac{1}{2} Y^R B_\mu$$

## Photon identification:

“rotation”:

$$\begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix} = \begin{pmatrix} c_W & s_W \\ -s_W & c_W \end{pmatrix} \begin{pmatrix} W_\mu^3 \\ B_\mu \end{pmatrix}, \quad c_W = \cos \theta_W, s_W = \sin \theta_W, \\ \theta_W = \text{mixing angle}$$

$$D_\mu^L \Big|_{A_\mu} = -\frac{i}{2} A_\mu \begin{pmatrix} -g_2 s_W - g_1 c_W Y^L & 0 \\ 0 & g_2 s_W - g_1 c_W Y^L \end{pmatrix} \stackrel{!}{=} ie A_\mu \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix}$$

- charged difference in doublet  $Q_1 - Q_2 = 1 \rightarrow g_2 = \frac{e}{s_W}$

- normalize  $Y^{L/R}$  such that  $g_1 = \frac{e}{c_W}$

$\hookrightarrow Y$  fixed by “Gell-Mann–Nishijima relation”:  $Q = T_I^3 + \frac{Y}{2}$



## Fermion–gauge-boson interaction:


$$\mathcal{L}_{\text{ferm, YM}} = \frac{e}{\sqrt{2}s_W} \overline{\Psi}_F^L \begin{pmatrix} 0 & W^+ \\ W^- & 0 \end{pmatrix} \Psi_F^L + \frac{e}{2c_W s_W} \overline{\Psi}_F^L \sigma^3 Z \Psi_F^L$$

$$- e \frac{s_W}{c_W} Q_f \overline{\psi}_f Z \psi_f - e Q_f \overline{\psi}_f A \psi_f \quad (f=\text{all fermions}, F=\text{all doublets})$$


## Feynman rules:



$$\frac{ie}{\sqrt{2}s_W} \gamma_\mu \omega_-$$



$$-iQ_f e \gamma_\mu$$



$$ie \gamma_\mu (g_f^+ \omega_+ + g_f^- \omega_-) = ie \gamma_\mu (v_f - a_f \gamma_5)$$

$$\text{with } g_f^+ = -\frac{s_W}{c_W} Q_f, \quad g_f^- = -\frac{s_W}{c_W} Q_f + \frac{T_{I,f}^3}{c_W s_W},$$

$$v_f = -\frac{s_W}{c_W} Q_f + \frac{T_{I,f}^3}{2c_W s_W}, \quad a_f = \frac{T_{I,f}^3}{2c_W s_W}$$

## gauge field Lagrangian (Yang-Mills Lagrangian)

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4}W_{\mu\nu}^a W^{a,\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu}$$

Field-strength tensors:


$$W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g_2 \epsilon^{abc} W_\mu^b W_\nu^c, \quad B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$

Lagrangian in terms of “physical” fields:

$$\begin{aligned} \mathcal{L}_{\text{YM}} = & -\frac{1}{2}(\partial_\mu W_\nu^+ - \partial_\nu W_\mu^+)(\partial^\mu W^{-,\nu} - \partial^\nu W^{-,\mu}) \\ & - \frac{1}{4}(\partial_\mu Z_\nu - \partial_\nu Z_\mu)(\partial^\mu Z^\nu - \partial^\nu Z^\mu) - \frac{1}{4}(\partial_\mu A_\nu - \partial_\nu A_\mu)(\partial^\mu A^\nu - \partial^\nu A^\mu) \\ & + \text{(trilinear interaction terms involving } AW^+W^-, ZW^+W^-) \\ & + \text{(quadrilinear interaction terms involving} \\ & \quad AAW^+W^-, AZW^+W^-, ZZW^+W^-, W^+W^-W^+W^-) \end{aligned}$$


## Feynman rules for gauge-boson self-interactions:

(fields and momenta incoming)



$$ieC_{WWV} \left[ g_{\mu\nu}(k_+ - k_-)_\rho + g_{\nu\rho}(k_- - k_V)_\mu + g_{\rho\mu}(k_V - k_+)_\nu \right]$$

with  $C_{WW\gamma} = 1$ ,  $C_{WWZ} = -\frac{c_W}{s_W}$



$$ie^2C_{WWVV'} \left[ 2g_{\mu\nu}g_{\rho\sigma} - g_{\mu\rho}g_{\sigma\nu} - g_{\mu\sigma}g_{\nu\rho} \right]$$

with  $C_{WW\gamma\gamma} = -1$ ,  $C_{WW\gamma Z} = \frac{c_W}{s_W}$ ,

$$C_{WWZZ} = -\frac{c_W^2}{s_W^2}, \quad C_{WWWW} = \frac{1}{s_W^2}$$

# Higgs mechanism $\Rightarrow$ masses of W and Z bosons

spontaneous breaking  $SU(2)_I \times U(1)_Y \rightarrow U(1)_Q$   
unbroken em. gauge symmetry, massless photon


Minimal scalar sector with complex scalar doublet  $\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ ,  $Y_\Phi = 1$

Scalar self-interaction via Higgs potential:

$$V(\Phi) = -\mu^2 \Phi^\dagger \Phi + \frac{\lambda}{4} (\Phi^\dagger \Phi)^2, \quad \mu^2, \lambda > 0,$$

=  $SU(2)_I \times U(1)_Y$  symmetric

$$V(\Phi) = \text{minimal for } |\Phi| = \sqrt{\frac{2\mu^2}{\lambda}} \equiv \frac{v}{\sqrt{2}} > 0$$



ground state  $\Phi_0$  (=vacuum expectation value of  $\Phi$ ) not unique

specific choice  $\Phi_0 = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix}$  not gauge invariant  $\Rightarrow$  spontaneous symmetry breaking

emg. gauge invariance unbroken, since  $Q\Phi_0 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \Phi_0 = 0$

Field excitations in  $\Phi$ :

$$\Phi(x) = \begin{pmatrix} \phi^+(x) \\ \frac{1}{\sqrt{2}} \left( v + H(x) + i\chi(x) \right) \end{pmatrix}$$


Gauge-invariant Lagrangian of Higgs sector:  $(\phi^- = (\phi^+)^\dagger)$

$$\mathcal{L}_H = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi) \quad \text{with } D_\mu = \partial_\mu - ig_2 \frac{\sigma^a}{2} W_\mu^a + i \frac{g_1}{2} B_\mu$$

$$= (\partial_\mu \phi^+) (\partial^\mu \phi^-) - \frac{iev}{2s_W} (W_\mu^+ \partial^\mu \phi^- - W_\mu^- \partial^\mu \phi^+) + \frac{e^2 v^2}{4s_W^2} W_\mu^+ W^{-,\mu}$$

$$+ \frac{1}{2} (\partial\chi)^2 + \frac{ev}{2c_W s_W} Z_\mu \partial^\mu \chi + \frac{e^2 v^2}{4c_W^2 s_W^2} Z^2 + \frac{1}{2} (\partial H)^2 - \mu^2 H^2$$

+ (trilinear  $SSS$ ,  $SSV$ ,  $SVV$  interactions)



+ (quadrilinear  $SSSS$ ,  $SSVV$  interactions)



Implications:

- gauge-boson masses:  $M_W = \frac{ev}{2s_W}$ ,  $M_Z = \frac{ev}{2c_W s_W} = \frac{M_W}{c_W}$ ,  $M_\gamma = 0$
- physical Higgs boson  $H$ :  $M_H = \sqrt{2\mu^2}$  = free parameter
- would-be Goldstone bosons  $\phi^\pm, \chi$ : unphysical degrees of freedom

general gauge: Goldstone fields  $\phi^\pm, \chi$  are present

required: gauge fixing term  $\mathcal{L}_{fix}$

$R_\xi$  gauge:


$$\mathcal{L}_{fix} = -\frac{1}{2\xi_\gamma} (F^\gamma)^2 - \frac{1}{2\xi_Z} (F^Z)^2 - \frac{1}{2\xi_W} (F^\pm)^2$$

with the gauge-fixing functionals  $F^a$ : ( $\xi_V$  = arbitrary gauge-fixing parameters)


$$F^\pm = \partial W^\pm \mp i\xi_W M_W \phi^\pm, \quad F^Z = \partial Z - \xi_Z M_Z \chi, \quad F^\gamma = \partial A$$

(notation:  $\partial A = \partial_\mu A^\mu, \dots$ )

- elimination of mixing terms  $(W_\mu^\pm \partial^\mu \phi^\mp), (Z_\mu \partial^\mu \chi)$  in Lagrangian  
 $\hookrightarrow$  decoupling of gauge and would-be Goldstone fields (no mix propagators)
- boson propagators:



$$D_{\mu\nu}^{VV}(k) = -i \left[ \frac{g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}}{k^2 - M_V^2} + \frac{k_\mu k_\nu}{k^2} \frac{\xi_V}{k^2 - \xi_V M_V^2} \right], \quad V = W, Z, \gamma$$



$$D^{SS}(k) = \frac{i}{k^2 - \xi_V M_V^2}, \quad S = \phi, \chi$$

- important special cases:

- ◇  $\xi_V = 1$ : ‘t Hooft–Feynman gauge  
 $\hookrightarrow$  convenient gauge-boson propagators  $\frac{-ig_{\mu\nu}}{k^2 - M_V^2}$
- ◇  $\xi_W, \xi_Z \rightarrow \infty$ : “unitary gauge”  
 $\hookrightarrow$  elimination of would-be Goldstone bosons

## Fermion masses

fermions in chiral representations of gauge symmetry

$$\begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \quad e_R \Rightarrow \text{mass term } m_e(\bar{e}_L e_R + \bar{e}_R e_L) = m_e \bar{e}e$$

not gauge invariant

**solution of the SM:** introduce Yukawa interaction

= new interaction of fermions with the Higgs field

gauge invariant interaction,  $g_e$  = Yukawa coupling constant

$$\mathcal{L}_{\text{Yuk}} = g_e [\bar{\psi}^L \Phi e_R + \bar{e}_R \Phi^\dagger \psi^L]$$



most transparent in unitary gauge

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \rightarrow \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H \end{pmatrix}$$

apply to the first lepton generation  $\psi^L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$ ,  $e_R$  :

$$\frac{g_e}{\sqrt{2}} \left[ (\overline{\nu}_L, \overline{e}_L) \begin{pmatrix} 0 \\ v + H \end{pmatrix} e_R + \overline{e}_R (0, v + H) \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \right]$$

$$= \underbrace{\frac{g_e}{\sqrt{2}} v}_{m_e} [\overline{e}_L e_R + \overline{e}_R e_L] + \underbrace{\frac{g_e}{\sqrt{2}} H}_{m_e/v} [\overline{e}_L e_R + \overline{e}_R e_L]$$

$$= m_e \bar{e}e + \frac{m_e}{v} H \bar{e}e$$

## 3 generations of leptons and quarks

Lagrangian for Yukawa couplings:

$$\mathcal{L}_{\text{Yuk}} = -\overline{\Psi}_L^L G_l \psi_l^R \Phi - \overline{\Psi}_Q^L G_u \psi_u^R \tilde{\Phi} - \overline{\Psi}_Q^L G_d \psi_d^R \Phi + \text{h.c.}$$

- $G_l, G_u, G_d = 3 \times 3$  matrices in 3-dim. space of generations ( $\nu$  masses ignored)
- $\tilde{\Phi} = i\sigma^2 \Phi^* = \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix}$  = charge conjugate Higgs doublet,  $Y_{\tilde{\Phi}} = -1$

Fermion mass terms:

mass terms = bilinear terms in  $\mathcal{L}_{\text{Yuk}}$ , obtained by setting  $\Phi \rightarrow \Phi_0$ :

$$\mathcal{L}_{m_f} = -\frac{v}{\sqrt{2}} \overline{\psi}_l^L G_l \psi_l^R - \frac{v}{\sqrt{2}} \overline{\psi}_u^L G_u \psi_u^R - \frac{v}{\sqrt{2}} \overline{\psi}_d^L G_d \psi_d^R + \text{h.c.}$$

↪ diagonalization by unitary field transformations ( $f = l, u, d$ )

$$\hat{\psi}_f^{L/R} \equiv U_f^{L/R} \psi_f^{L/R} \quad \text{such that} \quad \frac{v}{\sqrt{2}} U_f^L G_f (U_f^R)^\dagger = \text{diag}(m_f)$$

$$\Rightarrow \text{standard form:} \quad \mathcal{L}_{m_f} = -m_f \overline{\hat{\psi}}_f^L \hat{\psi}_f^R + \text{h.c.} = -m_f \overline{\hat{\psi}}_f \hat{\psi}_f$$

## Quark mixing:

- $\psi_f$  correspond to eigenstates of the gauge interaction
- $\hat{\psi}_f$  correspond to mass eigenstates,  
for **massless neutrinos** define  $\hat{\psi}_\nu^L \equiv U_l^L \psi_\nu^L \rightarrow$  **no lepton-flavour changing**

## Yukawa and gauge interactions in terms of mass eigenstates:

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = & -\frac{\sqrt{2}m_l}{v} \left( \phi^+ \overline{\hat{\psi}_{\nu_l}^L} \hat{\psi}_l^R + \phi^- \overline{\hat{\psi}_l^R} \hat{\psi}_{\nu_l}^L \right) + \frac{\sqrt{2}m_u}{v} \left( \phi^+ \overline{\hat{\psi}_u^R} V \hat{\psi}_d^L + \phi^- \overline{\hat{\psi}_d^L} V^\dagger \hat{\psi}_u^R \right) \\ & - \frac{\sqrt{2}m_d}{v} \left( \phi^+ \overline{\hat{\psi}_u^L} V \hat{\psi}_d^R + \phi^- \overline{\hat{\psi}_d^R} V^\dagger \hat{\psi}_u^L \right) - \frac{m_f}{v} i \text{sgn}(T_{I,f}^3) \chi \overline{\hat{\psi}_f} \gamma_5 \hat{\psi}_f \\ & - \frac{m_f}{v} (v + H) \overline{\hat{\psi}_f} \hat{\psi}_f, \end{aligned}$$

$$\begin{aligned} \mathcal{L}_{\text{ferm, YM}} = & \frac{e}{\sqrt{2}s_W} \overline{\hat{\Psi}_L^L} \begin{pmatrix} 0 & W^+ \\ W^- & 0 \end{pmatrix} \hat{\psi}_L^L + \frac{e}{\sqrt{2}s_W} \overline{\hat{\Psi}_Q^L} \begin{pmatrix} 0 & V W^+ \\ V^\dagger W^- & 0 \end{pmatrix} \hat{\psi}_Q^L \\ & + \frac{e}{2c_W s_W} \overline{\hat{\Psi}_F^L} \sigma^3 \not{Z} \hat{\Psi}_F^L - e \frac{s_W}{c_W} Q_f \overline{\hat{\psi}_f} \not{Z} \hat{\psi}_f - e Q_f \overline{\hat{\psi}_f} \not{A} \hat{\psi}_f \end{aligned}$$

- only charged-current coupling of quarks modified by  $V = U_u^L (U_d^L)^\dagger =$  **unitary**  
(Cabibbo–Kobayashi–Maskawa (CKM) matrix)
- **Higgs–fermion coupling strength**  $= \frac{m_f}{v}$

## Features of the CKM mixing:


- $V = 3$ -dim. generalization of Cabibbo matrix  $U_C$
- $V$  is parametrized by 4 free parameters: 3 real angles, 1 complex phase  
↪ complex phase is the only source of CP violation in SM

counting:

$$\begin{aligned} & \left( \begin{array}{c} \text{\#real d.o.f.} \\ \text{in } V \end{array} \right) - \left( \begin{array}{c} \text{\#unitarity} \\ \text{relations} \end{array} \right) - \left( \begin{array}{c} \text{\#phase diffs. of} \\ u\text{-type quarks} \end{array} \right) - \left( \begin{array}{c} \text{\#phase diffs. of} \\ d\text{-type quarks} \end{array} \right) - \left( \begin{array}{c} \text{\#phase diff. between} \\ u\text{- and } d\text{-type quarks} \end{array} \right) \\ & = 18 - 9 - 2 - 2 - 1 = 4 \end{aligned}$$

- no flavour-changing neutral currents in lowest order,  
flavour-changing suppressed by factors  $G_\mu(m_{q_1}^2 - m_{q_2}^2)$  in higher orders  
("Glashow–Iliopoulos–Maiani mechanism")

# The Standard Model Lagrangian



- renormalizable  $\Rightarrow$  precision calculations
- quantum effects in precision observables detectable
- involve Higgs mass dependence

## **6. Phenomenology of W and Z bosons and precision tests**

## Cross sections and decay widths

scattering process:  $a + b \rightarrow b_1 + b_2 + \dots + b_n$

$$|a(p_a), b(p_b)\rangle = |i\rangle, \quad |b_1(p_1), \dots, b_n(p_n)\rangle = |f\rangle$$

matrix element = probability amplitude for  $i \rightarrow f$ :

$$S_{fi} = \langle f | S | i \rangle$$

for  $i \neq f$ :  $S_{fi} = (2\pi)^4 \delta^4(P_i - P_f) \mathcal{M}_{fi} \left[ \frac{1}{(2\pi)^{3/2}} \right]^{n+2}$

$$P_i = p_a + p_b = P_f = p_1 + \dots + p_n \quad \text{momentum conservation}$$

factors  $(2\pi)^{-3/2}$  from wave function normalization  
(plane waves)

$\mathcal{M}_{fi}$  from Feynman graphs and rules

probability for scattering into phase space element  $d\Phi$ :

$$dW_{fi} = |S_{fi}|^2 d\Phi, \quad d\Phi = \frac{d^3 p_1}{2p_1^0} \cdots \frac{d^3 p_n}{2p_n^0}$$

$$\frac{d^3 p_i}{2p_i^0} = d^4 p_i \delta(p_i^2 - m_i^2) \quad \text{Lorentz invariant phase space}$$

differential cross section:

$$d\sigma = \frac{(2\pi)^4}{4\sqrt{(p_a \cdot p_b)^2 - m_a^2 m_b^2}} |\mathcal{M}_{fi}|^2 (2\pi)^{-3n} \delta^4(P_i - P_f) \frac{d^3 p_1}{2p_1^0} \cdots \frac{d^3 p_n}{2p_n^0}$$



decay process:  $a \rightarrow b_1 + b_2 + \dots + b_n$

$$|a(p_a)\rangle = |i\rangle, \quad |b_1(p_1), \dots, b_n(p_n)\rangle = |f\rangle$$

$$S_{fi} = (2\pi)^4 \delta^4(p_a - P_f) \mathcal{M}_{fi} \left[ \frac{1}{(2\pi)^{3/2}} \right]^{n+1}$$

decay width (differential):

$$d\Gamma = \frac{(2\pi)^4}{2m_a} |\mathcal{M}_{fi}|^2 (2\pi)^{-3n} \delta^4(p_a - P_f) \frac{d^3p_1}{2p_1^0} \dots \frac{d^3p_n}{2p_n^0}$$

## special case: 2-particle phase space

$$a + b \rightarrow b_1 + b_2, \quad a \rightarrow b_1 + b_2$$

### ● cross section

in the CMS,  $\vec{p}_a + \vec{p}_b = 0 = \vec{p}_1 + \vec{p}_2$

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2 s} \frac{|\vec{p}_1|}{|\vec{p}_a|} |\mathcal{M}_{fi}|^2$$

$$d\Omega = d\cos\theta d\phi, \quad \theta = \angle(\vec{p}_a, \vec{p}_1)$$

$$s = (p_a + p_b)^2 = E_{\text{CMS}}^2$$

### ● decay rate

for final state masses  $m_1 = m_2 = m$

$$\frac{d\Gamma}{d\Omega} = \frac{1}{64\pi^2 m_a} \sqrt{1 - \frac{4m^2}{m_a^2}} |\mathcal{M}_{fi}|^2$$

# features of the ew Standard Model

- Higgs boson probably found, all other particles confirmed
- consistent quantum field theory
  - in accordance with unitarity
  - renormalizable  $\Rightarrow$  predictions at higher orders
- formal parameters:  $g_2, g_1, v, \lambda, g_f, V_{\text{CKM}}$   
physical parameters:  $\alpha, M_W, M_Z, M_H, m_f, V_{\text{CKM}}$

# Basic parameters and relations

ew mixing angle:  $s_W \equiv \sin \theta_W, \quad c_W \equiv \cos \theta_W$

gauge coupling constants:  $g_2 = \frac{e}{s_W}, \quad g_1 = \frac{e}{c_W}$

vector boson masses:  $M_W = \frac{1}{2}g_2v = \frac{ev}{2s_W}$

$$M_Z = \frac{ev}{2s_W c_W} = \frac{M_W}{c_W}$$


$$s_W^2 = 1 - \frac{M_W^2}{M_Z^2}$$

neutral current (NC) couplings:  $a_f = \frac{g_2}{2c_W} T_3^f$

$$v_f = \frac{g_2}{2c_W} (T_3^f - 2Q_f s_W)$$

# observables and experiments

- Muon decay:




$$\mu^- \rightarrow \nu_\mu e^- \bar{\nu}_e$$

determination of the Fermi constant

$$G_\mu = \frac{\pi \alpha M_Z^2}{\sqrt{2} M_W^2 (M_Z^2 - M_W^2)} + \dots$$

- Z production (LEP1/SLC):




$$e^+ e^- \rightarrow Z \rightarrow f \bar{f}$$

various precision measurements at the Z resonance:  $M_Z, \Gamma_Z, \sigma_{\text{had}}, A_{\text{FB}}, A_{\text{LR}}, \text{etc.}$

⇒ good knowledge of the  $Z f \bar{f}$  sector


- W-pair production (LEP2/ILC):  $e^+ e^- \rightarrow WW \rightarrow 4f (+\gamma)$



– measurement of  $M_W$


–  $\gamma WW/ZWW$  couplings

– quartic couplings:  $\gamma\gamma WW, \gamma ZWW$




## experiments at hadron colliders

- **W production** (Tevatron/LHC):  $pp, p\bar{p} \rightarrow W \rightarrow l\nu_l(+\gamma)$




- measurement of  $M_W$
- bounds on  $\gamma WW$  coupling

- **top-quark production** (Tevatron/LHC):  $pp, p\bar{p} \rightarrow t\bar{t} \rightarrow 6f$



- measurement of  $m_t$

## μ decay



$$\mathcal{M} = \left( \frac{ig_2}{2\sqrt{2}} \right)^2 J_\rho^{(\mu)} \frac{-ig^{\rho\sigma}}{q^2 - M_W^2} J_\sigma^{(e)}$$

$$|q|^2 \simeq m_\mu^2 \ll M_W^2 : \quad \mathcal{M} = -\frac{g_2^2}{8M_W^2} J_\rho^{(\mu)} J^\rho(e)$$



**Fermi model** with point-like 4-fermion interaction:

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} J_\rho^{(\mu)} J^\rho(e) \quad \text{low-energy limit of SM}$$

$$\Rightarrow \boxed{\frac{G_F}{\sqrt{2}} = \frac{g_2^2}{8M_W^2} = \frac{e^2}{8s_W^2 c_W^2 M_Z^2} = \frac{\pi\alpha}{2s_W^2 c_W^2 M_Z^2} = \frac{\pi\alpha}{2(1 - M_W^2/M_Z^2)M_W^2}}$$

$$G_F = 1.16637(1) \cdot 10^{-5} \text{ GeV}^{-2}$$

# Z resonance



$$\mathcal{M} = J_{\mu}^{(e)} \frac{-ig^{\mu\nu}}{s - M_Z^2 + iM_Z\Gamma_Z} J_{\nu}^{(f)}$$

*propagator with finite width  $\Gamma_Z$  (unstable particle)*

$$\Gamma_Z = \sum_f \Gamma(Z \rightarrow f\bar{f}), \quad \Gamma(Z \rightarrow f\bar{f}) = \frac{M_Z}{12\pi} (v_f^2 + a_f^2)$$



*differential cross section at  $s = M_Z^2$ :*

$$\frac{d\sigma}{d\Omega} \sim (v_e^2 + a_e^2)(v_f^2 + a_f^2) (1 + \cos^2 \theta) + (2v_e a_e)(2v_f a_f) \cdot 2 \cos \theta$$

$$\Rightarrow \text{forward-backward asymmetry} \quad A_{FB} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B} = \frac{3}{4} A_e A_f$$

$$A_f = \frac{2v_f a_f}{v_f^2 + a_f^2}$$

*polarized cross section for  $e_{L,R}^-$ :*

$$\Rightarrow \text{left-right asymmetry} \quad A_{LR} = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R} = A_e$$

asymmetries determine  $\sin^2 \theta_W$

$$\frac{v_f}{a_f} = 1 - 4 |Q_f| \sin^2 \theta_W$$

# input from experiments


- **LEP1/SLC:**  $e^+e^- \rightarrow Z \rightarrow f\bar{f}$   
LEP1:  $\sim 4 \times 10^6$  events/experiment  
4 experiments (1989 – 1995)
- **LEP2:**  $e^+e^- \rightarrow W^+W^-$   
 $\mathcal{O}(10^4)$  W pairs (1996 – 2000)
- **Tevatron:**  $q\bar{q}' \rightarrow W \rightarrow l\nu, q\bar{q}'$   
(p $\bar{p}$ )  $q\bar{q}' \rightarrow t\bar{t}, t \rightarrow W^+b \rightarrow \dots$
- **low-energy experiments** ( $\mu$  decay,  $\nu N$  scattering,  $\nu e$  scattering, atomic parity violation, ... )

## experimental results (selection)

$M_Z$ [GeV]	$= 91.1875 \pm 0.0021$	0.002%
$\Gamma_Z$ [GeV]	$= 2.4952 \pm 0.0023$	0.09%
$\sin^2 \theta_{\text{eff}}^{\text{lept}}$	$= 0.23148 \pm 0.00017$	0.07%
$M_W$ [GeV]	$= 80.385 \pm 0.015$	0.02%
$m_t$ [GeV]	$= 173.2 \pm 0.9$	0.52%
$G_F$ [GeV <sup>-2</sup> ]	$= 1.16637(1)10^{-5}$	0.001%


loop effects are at least one order of magnitude larger than experimental uncertainties

# precise experiments need precise calculations




"I think you should be more explicit here in step two."



# example: 1-loop diagrams for $\mu$ decay amplitude






W self-energy




$Wl\nu_l$  vertex correction



box diagrams



Example of loop integral:



The diagram shows a loop integral with two external lines and a loop. The external lines are labeled with momentum  $p$  (in blue). The loop is a circle with two vertices. The top arc of the loop is labeled with momentum  $q$  (in red), and the bottom arc is labeled with momentum  $q + p$  (in red). Arrows on the loop indicate a clockwise direction.

$$\sim \int d^4 q \frac{1}{(q^2 - m_1^2) [(q + p)^2 - m_2^2]}$$

$$q \rightarrow \infty : \quad \sim \int^{\infty} \frac{q^3 dq}{q^4} = \int^{\infty} \frac{dq}{q} \rightarrow \infty$$

$\Rightarrow$  integral diverges for large  $q$

$\Rightarrow$  theory in this form not physically meaningful

- needs
- (i) regularization
  - (ii) renormalization

## Regularization:

theory modified such that expressions become mathematically meaningful

⇒ “regulator” introduced, removed at the end

e.g. cut-off in loop integral

$$\int_0^\infty d^4 q \rightarrow \int_0^\Lambda d^4 q; \quad \Lambda \rightarrow \infty \text{ at the end}$$

technically more convenient: dimensional regularization

$$\int d^4 q \rightarrow \int d^D q, \quad D = 4 - \varepsilon; \quad D \rightarrow 4 \text{ at the end}$$

## Renormalization:

- absorption of divergencies
- determination of physical meaning of parameters order by order in perturbation theory

add counterterms that absorb divergent parts

- parameters in  $\mathcal{L}$  are formal, “bare parameters”

$$g_0 = g + \delta g \text{ for a coupling,} \quad m_0 = m + \delta m \text{ for a mass}$$

- $g, m$  are “physical”, *i.e.* measurable



mass renormalization,  $m_0^2 = m^2 + \delta m^2$

Physical mass: pole of propagator


inverse propagator up to 1-loop order:

$$\begin{array}{ccccccc} \text{---} & + & \text{---} & \text{---} & + & \text{---} & + \dots \\ & & \text{---} \circ \text{---} & & & \times \text{---} & \\ p^2 - m^2 & & \Sigma(p^2) & & & -\delta m^2 & \end{array}$$

**on-shell renormalization:**  $\delta m^2 = \text{Re } \Sigma(m^2)$

charge renormalization:  $e_0 = e + \delta e$

$\delta e$  cancels loop contributions to  $ee\gamma$  vertex in the Thomson limit




$\Rightarrow e =$  elementary charge of classical electrodynamics

$$\text{fine-structure constant } \alpha(0) = \frac{e^2}{4\pi} = 1/137.03599976$$

$\delta e$  contains photon vacuum polarization  $\Pi^\gamma(k^2 = 0)$  :

$$\Pi^\gamma(0) = \underbrace{\Pi^\gamma(0) - \Pi^\gamma(M_Z^2)}_{\text{non-perturbative}} + \underbrace{\Pi^\gamma(M_Z^2)}_{\text{perturbative}}$$

# photon vacuum polarization



$$\Pi^\gamma(M_Z^2) - \Pi^\gamma(0) \equiv \Delta\alpha \quad \rightarrow \quad \alpha(M_Z) = \frac{\alpha}{1 - \Delta\alpha} \simeq \frac{1}{129}$$

$$\Delta\alpha = \Delta\alpha_{\text{lept}} + \Delta\alpha_{\text{had}},$$

$$\Delta\alpha_{\text{lept}} = 0.031498 \quad (3 - \text{loop})$$

$$\Delta\alpha_{\text{had}} = 0.02750 \pm 0.00033$$


$$= 0.02757 \pm 0.00010$$

*arXiv:1010.4180*


$$\Delta\alpha_{\text{had}} = -\frac{\alpha}{3\pi} M_Z^2 \operatorname{Re} \int_{4m_\pi^2}^{\infty} ds' \frac{R_{\text{had}}(s')}{s'(s' - M_Z^2 - i\epsilon)}$$

$$R_{\text{had}} =$$

$$\frac{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow \mu^+\mu^-)}$$




# $M_W - M_Z$ correlation



$$\frac{G_F}{\sqrt{2}} = \frac{\pi\alpha}{2M_W^2 (1 - M_W^2/M_Z^2)}$$

# $M_W - M_Z$ correlation



$$\frac{G_F}{\sqrt{2}} = \frac{\pi\alpha}{2M_W^2 (1 - M_W^2/M_Z^2)}$$

$$M_W = 80.939 \pm 0.002 \text{ GeV} \quad \text{from} \quad G_F, \alpha, M_Z$$

$$M_W = 79.965 \pm 0.005 \text{ GeV} \quad \text{with} \quad \alpha \rightarrow \alpha(M_Z)$$

$$M_W = 80.385 \pm 0.015 \text{ GeV} \quad \text{exp.} \quad 37\sigma / 28\sigma$$

## with loop contributions

$$\frac{G_F}{\sqrt{2}} = \frac{\pi\alpha}{2M_W^2 (1 - M_W^2/M_Z^2)} \cdot (1 + \Delta r)$$

$\Delta r$  : quantum correction

$$\Delta r = \Delta r(m_t, M_H)$$

$$\Delta r = \Delta\alpha - \frac{c_W^2}{s_W^2} \Delta\rho + \dots$$

$$\Delta\rho \sim \frac{m_t^2}{M_W^2}$$


determines W mass

$$M_W = M_W(\alpha, G_F, M_Z, m_t, M_H)$$


complete at 2-loop order

## 1-loop examples


- top quark






- Higgs boson



- gauge-boson self-couplings




full structure of SM



*2-loop examples*



# effects of higher-order terms on $\Delta r$




variation of  $\Delta r$  by 0.001  $\Rightarrow \delta M_W = 18 \text{ MeV}$



3-loop ( $\Delta\rho$ )  $\Rightarrow \delta M_W = 12 \text{ MeV}$

present exp. error:  $\Delta M_W = 15 \text{ MeV}$  / **theo: 4 MeV**

# LEP Electroweak Working Group



# Z resonance




- effective  $Z$  boson couplings with higher-order  $\Delta g_{V,A}$


$$v_f \rightarrow g_V^f = v_f + \Delta g_V^f, \quad a_f \rightarrow g_A^f = a_f + \Delta g_A^f$$

- effective ew mixing angle (for  $f = e$ ):




$$\sin^2 \theta_{\text{eff}} = \frac{1}{4} \left( 1 - \text{Re} \frac{g_V^e}{g_A^e} \right) = \kappa \cdot \left( 1 - \frac{M_W^2}{M_Z^2} \right)$$




a)





b)



c)



d)



*2-loop examples for Z couplings*

complete 2-loop calculation available for  $\sin^2 \theta_{\text{eff}}$

## EW 2-loop calculations for $\Delta r$

*Freitas, Hollik, Walter, Weiglein*

*Awramik, Czakon*

*Onishchenko, Veretin*

## EW 2-loop calculations for $\sin^2 \theta_{\text{eff}}$

*Awramik, Czakon, Freitas, Weiglein*

*Awramik, Czakon, Freitas*

*Hollik, Meier, Uccirati*

## universal terms at 3- and 4-loops (EW and QCD)

*van der Bij, Chetyrkin, Faisst, Jikia, Seidensticker*

*Faisst, Kühn, Seidensticker, Veretin*

*Boughezal, Tausk, van der Bij*


*Schröder, Steinhauser*

*Chetyrkin, Faisst, Kühn*

*Chetyrkin, Faisst, Kühn, Maierhofer, Sturm*

*Boughezal, Czakon*


## importance of two-loop calculations



lowest order:  $\sin^2 \theta_W = 1 - \frac{M_W^2}{M_Z^2} = 0.22290 \pm 0.00029$


exp. value:  $\sin^2 \theta_{\text{eff}} = 0.23153 \pm 0.00016$

# Global analysis within the SM



before the top quark was discovered ( $< 1995$ ):


indirect mass determination  $\Rightarrow m_t = 178 \pm 8^{+17}_{-20}$  GeV





before the top quark was discovered ( $< 1995$ ):

indirect mass determination  $\Rightarrow m_t = 178 \pm 8^{+17}_{-20}$  GeV




top discovery: *Tevatron 1995*

$m_t = 180 \pm 12$  GeV

before the top quark was discovered ( $< 1995$ ):

indirect mass determination  $\Rightarrow m_t = 178 \pm 8^{+17}_{-20}$  GeV



top discovery: *Tevatron 1995*


$$m_t = 180 \pm 12 \text{ GeV}$$

today:


$$m_t = 173.2 \pm 0.9 \text{ GeV}$$

# The way to the Higgs boson

development of bounds from direct and indirect searches



# Global fit to the Higgs boson mass




blueband: Theory uncertainty

“Precision Calculations  
at the  $Z$  Resonance”

CERN 95-03

*[Bardin, WH, Passarino (eds.)]*

$M_H < 161 \text{ GeV}$  (at 95% C.L.)



after the 2011 results  
from the LHC  
on the Higgs boson mass

$$M_H < 152 \text{ GeV} \quad (95\% \text{ C.L.})$$

$$M_H = 94_{-24}^{+29} \text{ GeV}$$

## **7. Higgs bosons**

Higgs potential: 
$$V = -\mu^2 (\Phi^\dagger \Phi)^2 + \frac{\lambda}{4} (\Phi^\dagger \Phi)^4$$

Higgs field in unitary gauge: 
$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix}$$

$H(x)$  : *real scalar field, describes neutral spin-0 bosons*

minimum of  $V$ : 
$$v = \frac{2\mu}{\sqrt{\lambda}}, \quad M_H = \mu\sqrt{2}$$

$$\Rightarrow \lambda = \frac{4\mu^2}{v^2} = \frac{2M_H^2}{v^2}$$

$$V = \frac{M_H^2}{2} H^2 + \frac{M_H^2}{2v} H^3 + \frac{M_H^2}{8v^2} H^4$$

$M_H$  is the only free parameter

Higgs potential: 
$$V = -\mu^2 (\Phi^\dagger \Phi)^2 + \frac{\lambda}{4} (\Phi^\dagger \Phi)^4$$

Higgs field in unitary gauge: 
$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix}$$

$H(x)$  : *real scalar field, describes neutral spin-0 bosons*

minimum of  $V$ : 
$$v = \frac{2\mu}{\sqrt{\lambda}}, \quad M_H = \mu\sqrt{2}$$

$$\Rightarrow \lambda = \frac{4\mu^2}{v^2} = \frac{2M_H^2}{v^2}$$

$$V = \frac{M_H^2}{2} H^2 + \frac{M_H^2}{2v} H^3 + \frac{M_H^2}{8v^2} H^4$$

$M_H$  is the only free parameter


general gauge: 
$$\Phi(x) = \begin{pmatrix} \phi^+(x) \\ \frac{1}{\sqrt{2}} [v + H(x) + i\chi(x)] \end{pmatrix}$$



## gauge invariant Lagrangian of the Higgs sector

$$\begin{aligned}
 \mathcal{L}_H &= (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi) \quad \text{with } D_\mu = \partial_\mu - ig_2 \frac{\sigma^a}{2} W_\mu^a + i \frac{g_1}{2} B_\mu \\
 &= (\partial_\mu \phi^+) (\partial^\mu \phi^-) - \frac{iev}{2s_W} (W_\mu^+ \partial^\mu \phi^- - W_\mu^- \partial^\mu \phi^+) + \frac{e^2 v^2}{4s_W^2} W_\mu^+ W^{-,\mu} \\
 &\quad + \frac{1}{2} (\partial \chi)^2 + \frac{ev}{2c_W s_W} Z_\mu \partial^\mu \chi + \frac{e^2 v^2}{4c_W^2 s_W^2} Z^2 + \frac{1}{2} (\partial H)^2 - \mu^2 H^2
 \end{aligned}$$

+ (trilinear  $SSS$ ,  $SSV$ ,  $SVV$  interactions)



+ (quadrilinear  $SSSS$ ,  $SSVV$  interactions)




**$\Rightarrow$  H-V-V gauge interactions, V=W and Z**

## gauge invariant Lagrangian of the Higgs sector

$$\begin{aligned}
 \mathcal{L}_H &= (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi) \quad \text{with } D_\mu = \partial_\mu - ig_2 \frac{\sigma^a}{2} W_\mu^a + i \frac{g_1}{2} B_\mu \\
 &= (\partial_\mu \phi^+) (\partial^\mu \phi^-) - \frac{iev}{2s_W} (W_\mu^+ \partial^\mu \phi^- - W_\mu^- \partial^\mu \phi^+) + \frac{e^2 v^2}{4s_W^2} W_\mu^+ W^{-,\mu} \\
 &\quad + \frac{1}{2} (\partial \chi)^2 + \frac{ev}{2c_W s_W} Z_\mu \partial^\mu \chi + \frac{e^2 v^2}{4c_W^2 s_W^2} Z^2 + \frac{1}{2} (\partial H)^2 - \mu^2 H^2
 \end{aligned}$$

+ (trilinear  $SSS$ ,  $SSV$ ,  $SVV$  interactions)




+ (quadrilinear  $SSSS$ ,  $SSVV$  interactions)




**⇒ H-V-V gauge interactions, V=W and Z**

$$\mathcal{L}_{\text{Yuk}} = - \sum_f \left( m_f + \frac{m_f}{v} H \right) \bar{\psi}_f \psi_f + \dots (ff \chi, \phi^\pm)$$

**⇒ H-f-f Yukawa interactions**



$$g_2 M_W, \quad g_2 \frac{M_Z}{c_W}$$





$$\frac{m_f}{v} = \frac{g_2 m_f}{2M_W}$$

$$\Gamma(H \rightarrow f \bar{f}) = N_c \frac{G_F M_H}{4\pi\sqrt{2}} m_f^2, \quad N_C = 3 (1) \text{ for quarks (leptons)}$$


$$\Gamma(H \rightarrow VV) = \frac{G_F M_H^3}{8\pi\sqrt{2}} F(r) \begin{pmatrix} 1 \\ 2 \end{pmatrix}_Z, \quad r = \frac{M_V}{M_H}$$

# Higgs boson decay channels

branching ratios  $BR(H \rightarrow X) = \frac{\Gamma(H \rightarrow X)}{\Gamma(H \rightarrow \text{all})}$






loop-induced (rare) decays




# Higgs decays into 4 fermions

also below  $VV$  threshold with one or two  $V$  off-shell






$$H \rightarrow VV \rightarrow 4f$$

needs also background processes + h.o.



Bredenstein et al.  $\rightarrow$  PROPHECY

$$H \rightarrow ZZ \rightarrow l^+l^- l^+l^-$$




signal + background




# Higgs boson production

Higgs production at LEP:





*excluded*  $M_H < 114 \text{ GeV}$

Higgs production at the Tevatron:



# Higgs production at the LHC



*Handbook of Higgs Cross sections,  
arXiv:1101.0593, arXiv:1201.3084*



“ I think we have it . . . ”





“ I think we have it . . . ”



. . . which one?

# A Standard Model Higgs boson at the LHC?



**H mass ATLAS (GeV)**


**$125.5 \pm 0.2 \pm 0.6$**

**H mass CMS (GeV)**

**$125.7 \pm 0.3 \pm 0.3$**

**Theory:  $\sigma(pp \rightarrow H) \cdot BR(H \rightarrow X)$**

# cross section for Higgs-boson production – theory



**NLO:** Spira, Djouadi, Graudenz, Zerwas '91, '93  
 Dawson '91 ~80%

**NNLO:** RH, Kilgore '02  
 Anastasiou, Melnikov '02 ~30%  
 Ravindran, Smith, v. Neerven '03

**Resummation:**

Catani, de Florian, Grazzini, Nason '02  
 Ahrens, Becher, Neubert, Zhang '08 ~10%

**Electroweak:**

Actis, Passarino, Sturm, Uccirati '08  
 Aglietti, Bonciani, Degrassi, Vicini '04  
 Degrassi, Maltoni '04 ~5%  
 Djouadi, Gambino '94



**Mixed EW/QCD:**

Anastasiou, Boughezal, Petriello '09

**Fully differential NNLO:**


Anastasiou, Melnikov, Petriello '04  
 Catani, Grazzini '07

# Effective Theory:





$$\sigma_{\infty}^{\text{HO}} \equiv \sigma^{\text{LO}}(m_t) \left( \frac{\sigma^{\text{HO}}}{\sigma^{\text{LO}}} \right)_{m_t \rightarrow \infty}$$

# Moch, Vogt '05





# impact of electroweak contributions at NLO




# Theoretical bounds on Higgs boson mass

- perturbativity  $\rightarrow$  upper bound
- unitarity  $\rightarrow$  upper bound
- Landau pole  $\rightarrow$  upper bound
- vacuum stability  $\rightarrow$  lower bound

## perturbativity

decay widths into fermions:  $\Gamma(H \rightarrow f\bar{f}) = \Gamma_{\text{tree}} \cdot K_f$

decay widths into vector bosons:  $\Gamma(H \rightarrow V\bar{V}) = \Gamma_{\text{tree}} \cdot K_V$



[Ghinculov; Frinck, Kniehl, Riesselmann]

# unitarity

scattering of longitudinally polarized  $W$  bosons:

$$W_L W_L \rightarrow W_L W_L$$

$$\mathcal{M}_V =$$

The diagram shows the scattering amplitude  $\mathcal{M}_V$  for the process  $W_L W_L \rightarrow W_L W_L$ . It is composed of three terms:

- The first term is a t-channel exchange of a photon ( $\gamma$ ) or a Z boson. It consists of two incoming wavy lines labeled  $W$  on the left and two outgoing wavy lines labeled  $W$  on the right. A horizontal wavy line in the middle represents the exchange particle, with two vertices marked by black dots. The label  $\gamma, Z$  is placed below the exchange line.
- The second term is a s-channel exchange of a photon ( $\gamma$ ) or a Z boson. It consists of two incoming wavy lines on the left and two outgoing wavy lines on the right. A vertical wavy line in the middle represents the exchange particle, with two vertices marked by black dots. The label  $\gamma, Z$  is placed to the right of the exchange line.
- The third term is a four-point contact interaction. It consists of four wavy lines meeting at a central vertex marked by a black dot.

$$= -g^2 \frac{E^2}{M_W^2} + \mathcal{O}(1) \quad \text{for } E \rightarrow \infty$$

Extra contribution from scalar particle:

$$\mathcal{M}_S = \text{Diagram 1} + \text{Diagram 2}$$

$$= g_{WWH}^2 \frac{E^2}{M_W^4} + \mathcal{O}(1) \quad \text{for } E \rightarrow \infty$$

$$\mathcal{M} = \mathcal{M}_V + \mathcal{M}_S$$

⇒ terms with bad high-energy behavior cancel for

$$g_{WWH} = g M_W$$

for  $s \gg M_W^2$ , with  $t = -\frac{s}{2} (1 - \cos \theta)$ ,

$$\mathcal{M} \approx \frac{M_H^2}{v^2} \left( 2 + \frac{M_H^2}{s - M_H^2} + \frac{M_H^2}{t - M_H^2} \right)$$

partial wave expansion:

$$\mathcal{M}(s, t) = 8\pi \sum_{l=0}^{\infty} (2l + 1) P_l(\cos \theta) a_l$$

unitarity condition:  $|a_l| < 1$


project on  $l = 0$  partial wave:

$$\begin{aligned} a_0 &= \frac{1}{16\pi} \int_{-1}^1 d \cos \theta \mathcal{M}(s, t) \\ &= \frac{M_H^2}{8\pi v^2} \left[ 2 + \frac{M_H^2}{s - M_H^2} - \frac{M_H^2}{s} \log \left( 1 + \frac{s}{M_H^2} \right) \right] \\ &\approx \frac{M_H^2}{4\pi v^2} \quad \text{for } s \gg M_H^2 \end{aligned}$$

$$a_0 < 1 \quad \Rightarrow \quad M_H < 872 \text{ GeV}$$

# Landau pole

Higgs self coupling is scale dependent,  $\lambda(Q)$



variation with scale  $Q$  described by RGE

$$Q^2 \frac{d\lambda}{dQ^2} = \beta(\lambda) = \frac{3}{4\pi^2} \lambda^2$$

solution:

$$\lambda(Q) = \frac{\lambda(v)}{1 - \frac{3}{4\pi^2} \lambda(v) \log \frac{Q^2}{v^2}} \quad \text{with} \quad \lambda(v) = \frac{M_H^2}{2v^2}$$

diverges at scale  $Q = \Lambda_C$  (Landau pole)

$$\Lambda_C = v \exp\left(\frac{4\pi^2 v^2}{3M_H^2}\right)$$

self-coupling diverges at

$$\Lambda_C = v \exp\left(\frac{4\pi^2 v^2}{3M_H^2}\right)$$


maximum Higgs mass by condition  $\Lambda_C > M_H$

$$\Rightarrow M_H < 800 \text{ GeV}$$



# vacuum stability

top-quark Yukawa coupling  $g_t \sim m_t$  contributes to the running Higgs self coupling  $\lambda(Q)$  through top loop  $\sim g_t^4$



variation with scale  $Q$  described by RGE

$$Q^2 \frac{d\lambda}{dQ^2} = \frac{3}{4\pi^2} \left( \lambda^2 - \frac{m_t^4}{v^4} \right)$$

approximate solution:


$$\lambda(Q) = \lambda(v) - \frac{3m_t^4}{2\pi^2 v^4} \log \frac{Q}{v}$$



$$\lambda(Q) < 0 \quad \text{for} \quad Q > \Lambda_C \quad \rightarrow \text{vacuum not stable}$$

high value of  $\Lambda_C$  needs  $M_H$  large enough

combined effects:


$$\frac{d\lambda}{dt} = \frac{1}{16\pi^2} (12\lambda^2 - 3g_t^4 + 6\lambda g_t^2 + \dots)$$






[Degrassi et al. 2012]

# Status of the Standard Model



SM input now completely determined  $\Rightarrow$  PO uniquely predicted

	theo	exp
$\sin^2 \theta_{\text{eff}}$	$0.23152 \pm 0.00005 \pm 0.00005$	$0.23153 \pm 0.00016$
$M_W$ (GeV)	$80.361 \pm 0.006 \pm 0.004$	$80.385 \pm 0.015$



## few observables with not-so-good agreement

- in general, SM is in overall agreement with data
- yet a few quantities prefer to stand a bit apart ( $\sim 3\sigma$ )
  - the forward-backward asymmetry for b quarks,  $A_{\text{FB}}^{b\bar{b}}$  at the Z peak
  - the anomalous magnetic moment of the muon
  - the forward-backward asymmetry for top quarks at the Tevatron,  $p\bar{p} \rightarrow t\bar{t}$

no conclusive situation

# The success of the Standard Model

- impressive confirmation by a huge data sample from low to high energies, no significant deviations
- quantum effects have been established at many  $\sigma$
- perfect indirect and direct determination of the top quark
- now being repeated for the Higgs boson
- new particle around 126 GeV strong candidate for the Higgs boson
- if confirmed: Standard Model closed

Happy End of a successful story ?

# Shortcomings of SM

- no mass terms for neutrinos [introduce  $\nu_R$  ... ]
- hierarchy problem  $v \ll M_{\text{Pl}}, \quad M_H \ll M_{\text{Pl}}$
- large number of free parameters  $g_1, g_2, v, m_f, V_{\text{CKM}}$
- no further unification of forces
- missing link to gravity
  
- nature of dark matter?
- baryon asymmetry of the universe?



- next steps with upgraded LHC
  - confirm the Higgs boson properties
  - check versus electroweak precision measurements
  - or find deviations, new structures:
    - more Higgs bosons (doublets, singlet, .. )
    - supersymmetry (minimal or non-minimal)
    - new strong sector, substructure
    - ...



**extra slides**

## SM Higgs:

- $\lambda H^4$  term ad hoc
- Higgs boson mass: free parameter  $\sim \sqrt{\lambda}$
- no a-priori reason for a light Higgs boson

## SM Higgs:

- $\lambda H^4$  term ad hoc
- Higgs boson mass: free parameter  $\sim \sqrt{\lambda}$
- no a-priori reason for a light Higgs boson


SUSY Standard Model avoids these questions

$$H_2 = \begin{pmatrix} H_2^+ \\ v_2 + H_2^0 \end{pmatrix}, \quad H_1 = \begin{pmatrix} v_1 + H_1^0 \\ H_1^- \end{pmatrix}$$


couples to  $u$                       couples to  $d$

- SUSY gauge interaction  $\rightarrow H^4$  terms
- self coupling remains weak


# spectrum of Higgs bosons in the MSSM: $h^0$ , $H^0$ , $A^0$ , $H^\pm$



# spectrum of Higgs bosons in the MSSM: $h^0$ , $H^0$ , $A^0$ , $H^\pm$




$m_h^0$  strongly influenced by quantum effects, e.g.  $t$ ,  $\tilde{t}$



# sensitivity to mass/mixing parameters

$m_{h^0}$  prediction at different levels of accuracy:





$\tan \beta = 3, \quad M_{\tilde{Q}} = M_A = 1 \text{ TeV}, \quad m_{\tilde{g}} = 800 \text{ GeV}$

$X_t$  : top-squark mixing parameter

$$X_t = A_t - \mu \cot \beta$$



# allowed region for top-squark mass and mixing



[Heinemeyer, Staal, Weiglein '12]

compatible with light top-squarks  
ongoing experimental search