


Teilchenphysik mit kosmischen und mit erdgebundenen Beschleunigern



8. Cosmic Rays II


15.06.2015




Neutral Cosmic Rays



Electromagnetic Radiation



Electromagnetic Radiation & The Atmosphere



Emission of Electromagnetic Radiation

- Thermal Radiation:
 - for example the sun: Emits at 5700 K (surface temperature of the sun)
 - Two-particle interactions lead to thermal equilibrium

$$\propto e^{-\frac{E}{kT}}$$


- Non-thermal radiation:
 - low density of plasma particles
 - a few particles can reach very high energies in interactions

$$\propto E^{-\gamma}$$

- Thermal and non-thermal components of a gas can exist in parallel and can interact with each other



Cosmic Background



1. Radio background, synchrotron radiation of e^- in galactic B field
2. Microwave background, 2.7 K
3. Emissions of cold interstellar dust
4. Emissions of distant galaxies
5. Hot interstellar dust
6. Optical background: scattering of sunlight on interstellar dust
7. Ionised intergalactic medium
8. X-ray background
9. Diffuse gamma background

Annu. Rev. Astron. Astrophys. 9, 89 (1991)

Beobachtung bei verschiedenen Wellenlängen


Radio (10^{-6} eV)

IR (10^{-2} eV)

sichtbar (1 eV)

X-Ray (10^3 eV)

Gamma (10^9 eV)



- Central regions of the Milky Way hidden by dust

Beobachtung bei verschiedenen Wellenlängen


Radio (10^{-6} eV)

IR (10^{-2} eV)

sichtbar (1 eV)

X-Ray (10^3 eV)

Gamma (10^9 eV)



Beobachtung bei verschiedenen Wellenlängen


Radio (10^{-6} eV)

IR (10^{-2} eV)

sichtbar (1 eV)

X-Ray (10^3 eV)

Gamma (10^9 eV)



- Dust in the Milky Way transparent to IR: Observation of the galactic center

Beobachtung bei verschiedenen Wellenlängen


Radio (10^{-6} eV)

IR (10^{-2} eV)

sichtbar (1 eV)

X-Ray (10^3 eV)

Gamma (10^9 eV)



Beobachtung bei verschiedenen Wellenlängen


Radio (10^{-6} eV)

IR (10^{-2} eV)

sichtbar (1 eV)


X-Ray (10^3 eV)

Gamma (10^9 eV)



- Combination of all wavelengths provides a detailed picture

An Example: Cen A



- Combined image of Centaurus A, an active galaxy at a distance of 10 Mlyr
 - visible light: white
 - sub-mm: orange
 - x-ray: blue

Credit: X-ray: NASA/CXC/CfA/R.Kraft et al.; Submillimeter: MPIfR/ESO/APEX/A.Weiss et al.; Optical: ESO/WFI




Photonen: “Acceleration”

- Photons themselves are not accelerated since they are uncharged
 - ▶ The energy originates from charged particles
 - ▶ Acceleration of charged particles in shock fronts
 - ▶ Conversion to photons through various processes
- ▶ The photons only receive a fraction of the energy of the primary particle
- ▶ Photons are substantially suppressed compared to hadrons at the same energy
- In the TeV region:
 $\text{Flux}(\gamma) \sim 10^{-4} \times \text{Flux}(\text{hadrons})$
- ▶ A challenge for experiments: Excellent photon/hadron separation required



Photon Production


- Accelerated charges emit radiation
- On circular orbits in magnetic fields:
Synchrotron radiation
- Acceleration (deceleration!) in strong electric fields
of nuclei:
Bremsstrahlung



Photon Production


- Accelerated charges emit radiation
- On circular orbits in magnetic fields:
Synchrotron radiation
- Acceleration (deceleration!) in strong electric fields
of nuclei:
Bremsstrahlung

In addition: Thermal radiation!





Photon Production II

- Hadronic production of photons via meson production in hadronic interactions and consecutive decay




- Energy increase of a photon via scattering off highly energetic electrons:
(inverse Compton scattering)




Expected Gamma Ray Spectrum of Sources

- Depends on the mechanism of γ creation




Expected Gamma Ray Spectrum of Sources

- Depends on the mechanism of γ creation




Expected Gamma Ray Spectrum of Sources


- Depends on the mechanism of γ creation




Gamma Creation: The General Idea



Direct Measurement: Satellites: Chandra




Direct Measurement: Satellites: Fermi




- Satellites can cover the full sky
- Good resolution at relatively low energies
- The newest instrument:
Fermi (formerly GLAST (Gamma-ray Large Area Telescope))
 - started on 11.06.2008

The Fermi Sky



Photon and Hadron - induced Air Showers


100 GeV photon



100 GeV proton


100 GeV photon

100 GeV proton




- Separation based on shower shape:
 - hadron showers are substantially more spread out and more “uneven”

Airshower Cherenkov Telescopes




- Cherenkov light is created by electrons in showers in an altitude of $\sim 10 \text{ km}$
- ▶ The photons are spread over an area with a radius of $\sim 120 \text{ m}$ on ground level
- ▶ Detection with a telescope is possible within this area.


Large IACTs



MAGIC: A large Cherenkov Telescope




- 17 m mirror diameter (240 m^2)
- ultralight construction (carbon fiber)
- Active mirror control: Correction of mechanical distortions




- Laser system for the alignment of the mirrors

MAGIC: Separation of Photons and Hadrons



SNR IC 443



blue: x-ray
green: radio
red: optical


Neutron star

Supernova
remnant, 3000 to
30 000 years old,
distance 1.5 kpc

Credit: Chandra X-ray: NASA/CXC/B.Gaensler et al; ROSAT X-ray: NASA/ROSAT/Asaoka & Aschenbach; Radio Wide: NRC/DRAO/D.Leahy; Radio Detail: NRAO/VLA; Optical: DSS



SNR IC 443 bei hochenergetischen Gammas




Quelle von TeV
Gammas: MAGIC
und VERITAS
Hochenergetische
Elektronen or
Hadronen, in einer
dichten Wolke
eingefangen?

>100 MeV Gammas,
Fermi LAT

Position of the
neutron star

Supernova Acceleration: Age & Energy

- Age dependence of cosmic accelerators



~ 1000 years needed
to reach peak energy
(see lecture 6)


Weakening of shock
wave and magnetic
fields with increasing
age

Active Galactic Nuclei: AGNs


- Supermassive black holes (10^7 - 10^{10} solar masses) in the center of galaxies
- Accretion of matter
 - depending on configuration a jet can be formed
 - approx 5% of all galaxies are active

Electromagnetic radiation from AGNs:

- Infrared to X-Ray
- Infrared
- Radio, Gamma (non-thermal)




Formation of Gammas in Jets




- Electrons and protons are accelerated in shock fronts
- Synchrotron radiation in magnetic fields
- Inverse Compton processes
- Proton induced cascades resulting in photons from neutral pion decay


AGN M87



X-Ray:




Gamma:




Blazars: Special AGNs

- A class of objects that exhibits strong gamma emission with pronounced time variations
 - first observation for BL Lacertae (BL Lac)
- AGN with a jet that points almost directly at earth
- Additional intensity gain due to relativistic effects
 - Time variation for example due to overtaking shock fronts that run through regions filled with matter, ...



AGNs: Photon Spectrum




- Typical “double hump” structure observed

Low Energy Cosmic Rays

- A complete particle physics detector in space
 - The goal: Search for antimatter in cosmic rays, detailed study of the composition, search for new phenomena

- A co
- Th
- se

AMS 02




AMS - Since 4 years on the ISS

- Successful start on May 16 2011 - Data taking since then




AMS - Illustration of Events




20 GeV electron

AMS - Illustration of Events




Composition of Cosmic Rays

- Comparison to the isotope abundance in the solar system (known from absorption lines in the sun, meteorites)
- Agreement for medium-mass nuclei: Maxima for even Z, A: stable nuclei preferred in fusion reactions in stars
- Deviation for light nuclei: Acceleration less efficient for H, He
- Li, Be, B are “burned” in stars instantly
 - Elements heavier than Li do not occur primordially
 - these are created in cosmic rays via spallation




Energy Dependence of Composition



Boron does not occur in primary cosmic rays: production via spallation,
Carbon is a primary component
⇒ Boron observed predominantly at low energy!

Energy Dependence of Composition



CREAM 2008



Both Oxygen and Carbon are primary particles in cosmic rays, both have $Z/A = 0.5$: identical energy spectrum!

Positrons in Cosmic Rays



- Interesting (and unexpected!) results by PAMELA (Nature, April 2009):



arXiv:0810.4995


Positrons in Cosmic Rays

- Interesting (and unexpected!) results by PAMELA (Nature, April 2009):



arXiv:0810.4995


Possible Explanation: Dark Matter



- Triggered a lot of activity: Several hundred papers with Dark Matter interpretations of PAMELA data within a few months


Would lead to a “Peak” in positrons, at higher energies the curve should go back to the expected behavior

Dark Matter: Consequence on Positron Fraction



Less Spectacular Explanation



- There has to be another primary source of positrons!
 - Production von positrons for example in near-by pulsars: Highly energetic gamma rays produce e^+e^- pairs



Electrons (and positrons) quickly loose energy in the ISM via Compton scattering and synchrotron radiation
⇒ Sources for highly-energetic positrons have to be close!

Atoyan et al., PRD 52, 3265 (1995)


Electrons and Positrons from Pulsars



Atoyan et al., PRD 52, 3265 (1995)

- ⇒ The PAMELA results can also be explained by a few (or a single) close pulsar, candidates do exist!


Electrons and Positrons from Pulsars



⇒ The PAMELA results can also be explained by a few (or a single) close pulsar, candidates do exist!

Confirmation by AMS and Fermi


Phys. Rev. Lett. 110, 141102 (2013)



- AMS proves: the effect is real
 - also supported by Fermi results
- Detailed analyses show (up to now) no anisotropy in the distribution of the positron excess - no preferred direction

Confirmation by AMS and Fermi

Phys. Rev. Lett. 110, 141102 (2013)



- AMS proves: the effect is real
 - also supported by Fermi results
- Detailed analyses show (up to now) no anisotropy in the distribution of the positron excess - no preferred direction

⇒ Consistent with “particle physics origin” (Dark Matter), but other explanations can not be excluded: Pulsars, possible mistakes in propagation models, ...

Summary

- Electromagnetic radiation is important: The combination of different spectral ranges allows detailed investigations of sources
- Highly energetic photons are observed with Cherenkov telescopes
 - Production via synchrotron radiation, inverse Compton processes and hadron decays
 - Provides insight into acceleration mechanisms
- Gamma sources in the Milky Way and beyond:
 - Pulsars
 - Active galactic nuclei, Blazars
- Composition of cosmic rays at low energy well understood - primary and secondary components
- Exciting observation: Positron excess A hint of New Physics, or pulsars or shortcomings of our understanding of particle propagation in the galaxy?



Summary

- Electromagnetic radiation is important: The combination of different spectral ranges allows detailed investigations of sources
- Highly energetic photons are observed with Cherenkov telescopes
 - Production via synchrotron radiation, inverse Compton processes and hadron decays
 - Provides insight into acceleration mechanisms
- Gamma sources in the Milky Way and beyond:
 - Pulsars
 - Active galactic nuclei, Blazars
- Composition of cosmic rays at low energy well understood - primary and secondary components
- Exciting observation: Positron excess A hint of New Physics, or pulsars or shortcomings of our understanding of particle propagation in the galaxy?

Next Lecture: 22.06., “Low-Energy Precision Experiments”, S. Bethke

Themenübersicht

13.04.	Einführung / Introduction
20.04.	Achtung - keine Vorlesung! No Lecture!
27.04.	Erdgebundene Beschleuniger / Accelerators
04.05.	Detektoren in der Nicht-Beschleuniger-Physik / Detectors
11.05.	Das Standardmodell / The Standard Model
18.05.	QCD und Jet Physik an Lepton Beschleunigern
25.05.	Pfingsten - Keine Vorlesung! No Lecture
01.06.	Kosmische Beschleuniger / Cosmic Accelerators
08.06.	Kosmische Strahlung I / Cosmic Rays I
15.06.	Kosmische Strahlung II / Cosmic Rays II
22.06.	Präzisionsexperimente (g-2) / Precision Experiments
29.06.	Neutrinos I
06.07.	Neutrinos II
13.07.	Dunkle Materie & Dunkle Energie / Dark Matter & Dark Energy

