

Hierarchical Soft Terms and Flavor Physics

Andrea Romanino
SISSA/ISAS

Giudice Nardecchia R, arXiv:0812.3610

SUSY flavour breaking sources

- New sources of flavour breaking (squarks only)

$$\begin{aligned} & \tilde{q}_L^\dagger \tilde{m}_{q_L}^2 \tilde{q}_L + \tilde{d}_R^\dagger \tilde{m}_{d_R}^2 \tilde{d}_R + \tilde{u}_R^\dagger \tilde{m}_{u_R}^2 \tilde{u}_R \\ & + \left(\tilde{d}_R^\dagger Y_D A_D \tilde{q}_L h_D + \tilde{u}_R^\dagger Y_U A_U \tilde{q}_L h_U + \text{h.c.} \right) \end{aligned}$$

- D-squark mass matrix (in the super-CKM basis)

$$\mathcal{M}_D^2 = \begin{pmatrix} LL & LR \\ RL & RR \end{pmatrix} = \mathcal{W}_D \mathcal{M}_D^{2 \text{diag}} \mathcal{W}_D^\dagger \quad (\tilde{d}_L^\dagger, \tilde{d}_R^\dagger) \mathcal{M}_D^2 \begin{pmatrix} \tilde{d}_L \\ \tilde{d}_R \end{pmatrix}$$

$$LL = \tilde{m}_{q_L}^2 + M_D^\dagger M_D + M_Z^2 z_D c_{2\beta} \mathbf{1}$$

$$RR = \tilde{m}_{d_R}^2 + M_D M_D^\dagger + M_Z^2 z_{D^c} c_{2\beta} \mathbf{1}$$

$$RL = -M_D (A_D + \mu \tan \beta)$$

$$M_D = \text{down quark mass matrix} = M_D^{\text{diag}}$$

Degeneracy and hierarchy

- About 100 physical parameters
- Large FCNC (and CPV) processes in most of the parameter space, (SUSY flavour problem)
- For $\tilde{m} < \text{TeV}$: need small 12/3 mixing + $\tilde{m}_d \approx \tilde{m}_s$ (barring alignment)
- Note: fermion masses also have peculiar structure: small 12/3 mixing + $m_d, m_s \ll m_b$; moreover, both correspond to an approximate U(2) symmetry
- U(3) badly broken by $Y_\dagger = O(1) \rightarrow$ expect $\tilde{m}_b \neq \tilde{m}_{d,s}$ How much?
Two opposite (complementary) limits:
 - Degeneracy: $\tilde{m}_b \approx \tilde{m}_s \approx \tilde{m}_d$
 - Hierarchy: $\tilde{m}_b \ll \tilde{m}_{s,d}$ (not incompatible with naturalness, see below)

Model-independent analysis

- Expand in small off-diagonal elements of squark mass matrix

$$\mathcal{M}^2 = \mathcal{M}_0^2 + \mathcal{M}_1^2$$

- **Degeneracy:** $\mathcal{M}_0^2 = \tilde{m}^2 \mathbf{1}$ [MIA, Gabbiani Gabrielli Masiero Silvestrini '96]

$$A(\Delta F = 1)_{ij} = x f^{(1)}(x) \delta_{ij}$$
$$A(\Delta F = 2)_{ij} = \frac{x^2}{3!} g^{(3)}(x) \delta_{ij}^2$$
$$\delta_{ij} \equiv \frac{(\mathcal{M}_1^2)_{ij}}{\tilde{m}^2}, \quad x \equiv \frac{\tilde{m}^2}{M^2}$$

- δ_{ij} : source of flavour violation, process independent
- f, g : loop functions (process dependent, flavour conserving)
- # of derivatives and factorial \leftrightarrow # of identical propagators (-1)
- OK in case of short RGE running, in any case useful recipe up to (harmless? see below) $O(1)$ differences between f and f'

Model-independent analysis

- Expand in small off-diagonal elements of squark mass matrix

$$\mathcal{M}^2 = \mathcal{M}_0^2 + \mathcal{M}_1^2$$

- Hierarchy:

[Cohen Kaplan Lepeintre Nelson '97]

$$\mathcal{M}_0^2 = \left(\begin{array}{c|c} \text{heavy} & \\ \hline & \tilde{m}^2 \\ \hline & \\ \hline & \text{heavy} \\ & \tilde{m}^2 \end{array} \right) \quad \begin{aligned} A(\Delta F = 1)_{ij} &= f(x) \hat{\delta}_{ij} \\ A(\Delta F = 2)_{ij} &= g^{(1)}(x) \hat{\delta}_{ij}^2 \end{aligned}$$

- $\hat{\delta}_{ij} \equiv \mathcal{W}_{i3} \mathcal{W}_{3j}^\dagger \quad x = \frac{\tilde{m}^2}{M^2} \quad \hat{\delta}_{a3} = \frac{\mathcal{M}_{a3}^2}{\tilde{m}_a^2} \quad (a = 1, 2)$

- $\hat{\delta}_{ab} \approx \hat{\delta}_{a3} \hat{\delta}_{3b} \quad (a, b, = 1, 2)$

- $\hat{\delta}_{a3}^{LR} \approx \hat{\delta}_{a3}^{LL} \hat{\delta}_{33}^{LR} \quad \hat{\delta}_{33}^{LR} = -\frac{m_b (A_{33}^D + \mu \tan \beta)}{\tilde{m}^2}$

4 complex parameters + 3rd family mixing

Degeneracy vs Hierarchy


- Different correlation between $\Delta F = 1$ and $\Delta F = 2$ in the two cases

$$\left. \frac{A(\Delta F = 2)}{[A(\Delta F = 1)]^2} \right|_{\text{degeneracy}} = \frac{1}{6} \frac{g^{(3)}}{g^{(1)}} \left(\frac{f}{f^{(1)}} \right)^2 \left. \frac{A(\Delta F = 2)}{[A(\Delta F = 1)]^2} \right|_{\text{hierarchy}}.$$

E.g.: $A(\Delta m_{B_s})$ vs $A(b \rightarrow s\gamma)$
LL insertion $\left. \frac{g^{(3)}}{g^{(1)}} \left(\frac{f}{f^{(1)}} \right)^2 \right|_{x=1} \approx \frac{25}{27}$

Theoretical considerations on the hierarchical case

- Complementary to degeneracy
- The naturalness bound on $\tilde{m}_{1,2}$ is milder than the one on \tilde{m}_3 [Dimopoulos Giudice '95]
- Effective supersymmetry [Cohen Kaplan Lepeintre Nelson '97]
- Alleviates the SUSY flavour problem: $\hat{\delta} \lesssim \tilde{m}/\tilde{m}_{1,2}$
- Related to physical quantities: squark masses, CKM angles



Bounds on δ 's

Hierarchy

Degeneracy

$D_0 - \bar{D}_0$ mixing

$$\left| \hat{\delta}_{ut}^{LL} \hat{\delta}_{ct}^{LL*} \right| < 8.0 \times 10^{-3} \left(\frac{m_{\tilde{t}}}{350 \text{ GeV}} \right) \quad \left| \delta_{uc}^{LL} \right| < 3.4 \times 10^{-2} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right)$$

$B \rightarrow X_s \gamma$

$$\begin{array}{|l} \left| \text{Re}(\hat{\delta}_{sb}^{LL}) \right| < 2.2 \times 10^{-2} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right)^2 \left(\frac{10}{\tan \beta} \right) \\ \left| \text{Im}(\hat{\delta}_{sb}^{LL}) \right| < 6.7 \times 10^{-2} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right)^2 \left(\frac{10}{\tan \beta} \right) \end{array} \quad \begin{array}{|l} \left| \text{Re}(\delta_{sb}^{LL}) \right| < 3.8 \times 10^{-2} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right)^2 \left(\frac{10}{\tan \beta} \right) \\ \left| \text{Im}(\delta_{sb}^{LL}) \right| < 1.1 \times 10^{-1} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right)^2 \left(\frac{10}{\tan \beta} \right) \end{array}$$

$$|\hat{\delta}_{sb}| \gtrsim 4 \cdot 10^{-2}$$

Δm_{B_s}

$$\begin{array}{|l} \left| \text{Re}(\hat{\delta}_{sb}^{LL}) \right| < 9.4 \times 10^{-2} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \\ \left| \text{Im}(\hat{\delta}_{sb}^{LL}) \right| < 7.2 \times 10^{-2} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \end{array} \quad \begin{array}{|l} \left| \text{Re}(\delta_{sb}^{LL}) \right| < 4.0 \times 10^{-1} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right) \\ \left| \text{Im}(\delta_{sb}^{LL}) \right| < 3.1 \times 10^{-1} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right) \end{array}$$

$$\begin{array}{l} \tilde{m} = M_3 = \mu \\ A = 0 \end{array}$$

$B_d^0 - \bar{B}_d^0$ mixing

$$\begin{array}{|l} \left| \text{Re}(\hat{\delta}_{db}^{LL}) \right| < 4.3 \times 10^{-3} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \\ \left| \text{Im}(\hat{\delta}_{db}^{LL}) \right| < 7.3 \times 10^{-3} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \end{array} \quad \begin{array}{|l} \left| \text{Re}(\delta_{db}^{LL}) \right| < 1.8 \times 10^{-2} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right) \\ \left| \text{Im}(\delta_{db}^{LL}) \right| < 3.1 \times 10^{-2} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right) \end{array}$$

$$|\hat{\delta}_{db}| \gtrsim 0.8 \cdot 10^{-2}$$

Δm_K


$$\sqrt{\left| \text{Re}(\hat{\delta}_{db}^{LL} \hat{\delta}_{sb}^{LL*})^2 \right|} < 1.0 \times 10^{-2} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \quad \sqrt{\left| \text{Re}(\delta_{ds}^{LL})^2 \right|} < 4.2 \times 10^{-2} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right)$$

$$|\hat{\delta}_{ds}| \gtrsim 3 \cdot 10^{-4}$$

ϵ_K

$$\sqrt{\left| \text{Im}(\hat{\delta}_{db}^{LL} \hat{\delta}_{sb}^{LL*})^2 \right|} < 4.4 \times 10^{-4} \left(\frac{m_{\tilde{b}}}{350 \text{ GeV}} \right) \quad \sqrt{\left| \text{Im}(\delta_{ds}^{LL})^2 \right|} < 1.8 \times 10^{-3} \left(\frac{m_{\tilde{q}}}{350 \text{ GeV}} \right)$$


$u \leftrightarrow c$ transitions ($D^0 - \bar{D}^0$)



68%, 95% C.L.

$$\tilde{m} = M_3 = 350 \text{ GeV}$$


$b \leftrightarrow d$ transitions ($\Delta m_{Bd}, \sin 2\beta_{\text{eff}}$)



68%, 95% C.L.

$\tilde{m} = M_3 = 350 \text{ GeV}$

$b \leftrightarrow s$ transitions ($\Delta m_{B_s}, b \rightarrow s\gamma, \phi_{B_s}$)




95% C.L.

$$\tilde{m} = M_3 = \mu = 350 \text{ GeV}, \tan\beta = 10, A = 0$$


ϕ_{B_s}

$$\langle B_s | H_{\text{eff}}^{\text{full}} | \bar{B}_s \rangle = C_{B_s} e^{2i\phi_{B_s}} \langle B_s | H_{\text{eff}}^{\text{SM}} | \bar{B}_s \rangle$$

$$\langle B_s | H_{\text{eff}}^{\text{SM}} | \bar{B}_s \rangle = A_s^{\text{SM}} e^{-2i\beta_s} \quad \beta_s = \arg(-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)) = 0.018 \pm 0.001$$



(*) Significance reduced from $\sigma \sim 3.0$ C_{B_s} to $\sigma \sim 2.5$ due to the correlation between $\Delta\Gamma$'s and ϕ_s in D0 likelihood



Summary

- $\tilde{m}_3 \ll \tilde{m}_{1,2}$ not incompatible with naturalness
- Welcome to alleviate the SUSY flavour problem
- Useful to describe scenarios where the mass separation is moderate but sufficient to make the degeneracy assumption a poor starting point. Which is somewhat expected because of $U(3)$ breaking by Y_+
- Complementary to the degeneracy assumption in a general study of SUSY flavour effects, with predictions different by factors $O(1-10)$
- Predictions
 - Less parameters than in the case of degeneracy: $d \leftrightarrow s$ via $d \leftrightarrow b \leftrightarrow s$
 $s_L \leftrightarrow b_R$ via $s_L \leftrightarrow b_L \leftrightarrow b_R$ (under assumptions)
 - Specific, distinct correlation $\Delta F = 1$ vs $\Delta F = 2$, with larger effects in $\Delta F = 2$ allowed
 - Experiment has begun to explore the range of expected $b \leftrightarrow s$ effects

Inputs

Parameter	Value	Gaussian (σ)	Uniform ($\frac{\Delta}{2}$)	Reference
$ \varepsilon_K $	2.229×10^{-3}	0.012×10^{-3}	—	[19]
Δm_K (ps $^{-1}$)	5.292×10^{-3}	0.009×10^{-3}	—	[19]
BR($B \rightarrow X_s \gamma$)	3.55×10^{-4}	0.26×10^{-4}	—	[20]
Δm_{B_s} (ps $^{-1}$)	17.77	0.12	—	[19]
Δm_{B_d} (ps $^{-1}$)	0.507	0.005	—	[19]
$ M_{12}^D $ (ps $^{-1}$)	7.7×10^{-3}	2.5×10^{-3}	—	[21]
$\bar{\rho}$	0.167	0.051	—	[22]
$\bar{\eta}$	0.386	0.035	—	[22]
λ	0.2255	0.010	—	[19]
$ V_{cb} $	41.2×10^{-3}	1.1×10^{-3}	—	[19]
F_K (GeV)	0.160	—	—	[19]
F_{B_d} (MeV)	189	27	—	[23]
$F_{B_s} \sqrt{B_s}$ (MeV)	262	35	—	[23]
F_D (MeV)	201	3	17	[24]
\hat{B}_K	0.79	0.04	0.08	[24]
B_1^B	0.88	0.04	0.10	[24]
η_{cc}	0.47	0.04	—	[25]
η_{ct}	0.5765	0.0065	—	[25]
η_{tt}	1.43	0.23	—	[25]
\bar{m}_t (GeV)	161.2	1.7	—	[24]
\bar{m}_b (GeV)	4.21	0.08	—	[24]
\bar{m}_c (GeV)	1.224	0.057	—	[26]