


The search for the (SM) Higgs Boson

- theoretical basics
- Higgs production and decay
- Higgs search in e^+e^- annihilation
 - direct search
 - indirect mass limits from electroweak radiation corrections
- Higgs searches in hadron collisions
 - Tevatron
 - LHC (-> next lecture)

The Standard Model Higgs Boson: theoretical basics and expectations

- gauge field theory with **gauge symmetry** in weak isospin/hyper charge [SU(2) \times U(1)] to describe electromagnetic and weak interactions of quarks and leptons:
includes **massless** gauge bosons (γ , Z^0 , W^+ , W^-) and fermions
- any attempt to include mass terms breaks gauge symmetry and destroys renormalizability of the theories
- Englert, Brout and Higgs (1964): **spontaneous symmetry breaking** (generates mass, keeps renormalizability):
- introduction of complex SU(2) doublets of scalar fields with a potential
of $V(\phi) = \lambda (\phi^+ \phi)^2 - \mu^2 \phi^+ \phi$; with $\lambda, \mu^2 > 0$; $\phi = \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}$
- V does not have minimum at $\phi = 0$, but at
 $|\phi| = \sqrt{\frac{\mu^2}{2\lambda}} = \frac{\nu}{\sqrt{2}}$
- 3 of the 4 real degrees of freedom are used to generate the longitudinal spin d.o.f. of Z^0 and W^\pm ;
4. d.o.f. \rightarrow physical Higgs particle!



theoretical basis and expectations

- inserting ϕ in Lagrange function results in 3 massive vector fields, 1 massless vector-field, plus one massive scalar field with



$$M_W = \frac{1}{2}gv \quad \Rightarrow \quad v = 246 \text{ GeV}$$

$$M_Z = M_W / \cos \theta_w \quad (g = e/\sin \theta_w)$$

$$M_\gamma = 0$$

$$M_H = 2\mu^2 = 2\lambda v^2$$

- introduction of Yukawa-couplings g_f between ϕ and the fermion fields: generates fermion masses $m_f = g_f v / \sqrt{2}$
- fundamental fermion-Higgs couplings:




theoretical basis and expectations

theoretical bounds for M_H from self-consistency arguments of the Standard-Model:

- upper bounds: perturbativity
- lower bounds: vacuum stability


n.b.: if SM is valid only up to
 $\Lambda = \mathcal{O}(1 \text{ TeV})$,
then $M_H = 50 \dots 1000 \text{ GeV}$

n.b.: if SM is valid up to $\Lambda = \mathcal{O}(M_{\text{Planck}})$
then $M_H \sim 125 \dots 180 \text{ GeV}$




Λ : energy scale up to which SM is valid


Higgs: production and decays




Higgs-radiation



W - (Z -) fusion




Gluon - Fusion




Higgs-radiation („associate production“)

Higgs-decay:

predominantly into heaviest,
kinematically accessible pair of
leptons or bosons



Higgs: decays




$M_H < \sim 135 \text{ GeV}$: dominanter Zerfallskanal $H \rightarrow b\bar{b}$


$M_H > \sim 135 \text{ GeV}$: dominanter Zerfallskanal $H \rightarrow W^+W^-$

Higgs: decays

Low mass region, e.g. $m_H = 125$ GeV:



High mass region, e.g. $m_H = 300$ GeV:



Ordered by the sensitivity to the signal:


- $H \rightarrow ZZ \rightarrow (\ell^-\ell^+)(\ell^+\ell^-)$
- $H \rightarrow \gamma\gamma$
- $H \rightarrow WW \rightarrow (\ell^+\nu)(\ell^-\nu)$
- $H \rightarrow \tau^+\tau^-$ (large background)
- $H \rightarrow b\bar{b}$ (large background)

Ordered by the sensitivity to the signal:

- $H \rightarrow ZZ \rightarrow (\ell^+\ell^-)(\ell^+\ell^-)$
- $H \rightarrow ZZ \rightarrow (\ell^+\ell^-)(\nu\nu)$
- $H \rightarrow ZZ \rightarrow (\ell^+\ell^-)(q\bar{q})$
- $H \rightarrow WW \rightarrow (\ell^-\nu)(\ell^-\nu)$
- $H \rightarrow WW \rightarrow (\ell^-\nu)(q\bar{q})$



Higgs: production

e^+e^- annihilation





Higgs: production

Standard Model Higgs Boson @ Tevatron



Higgs: production



Standard Model Higgs Boson @ LHC (7 TeV)




Higgs: production

Standard Model Higgs Boson @ LHC (14 TeV)


cross section (pb)



Higgs production cross-sections




Higgs search




Higgs-search in e^+e^- annihilation: direct

production:




decay channel ($e^+e^- \rightarrow HZ$):




51%

$WW \rightarrow qqqq, ZZ \rightarrow bbqq$
QCD 4jets




15%

$WW \rightarrow qq\nu\bar{\nu}, ZZ \rightarrow bb\nu\bar{\nu}$



2.4%

$WW \rightarrow qq\tau\nu, ZZ \rightarrow qq\tau\tau$
QCD (low-mult. jets)




5.1%

4.9%

$ZZ \rightarrow bbll$

search includes $\sim 80\%$ of all final states with $\sim 40 - 50\%$ selection efficiency

Higgs-search in e^+e^- annihilation: direct




L3: Candidate event
for $ee \rightarrow HZ \rightarrow ee\bar{q}\bar{q}$

LEP:
from direct search
 $M_H > 114.1 \text{ GeV}$

Higgs-search in e^+e^- annihilation: indirect


radiation corrections in SM:

photonic corrections:



corrections $\sim 100\%$, selection dependent;
factorisable: $(1 + \delta_{rad})$

non-photonic corrections:



corrections $\sim 10\%$, selection independent;
can be absorbed into running coupling constants:

- $\sin^2\theta_{eff}(s)$
- $\alpha(s) = \frac{\alpha}{1 - \Delta\alpha} ; \quad \Delta\alpha = 0.064 \text{ bei } \sqrt{s} = M_Z$
- $N_{c,f} \left(1 + \frac{\alpha_s}{\pi} + 1.4 \left(\frac{\alpha_s}{\pi} \right)^2 + \dots \right) \text{ (für Quarks)}$
- $\frac{M_W^2}{M_Z^2} = \rho \cdot \cos^2 \theta_w \quad \text{mit} \quad \rho = \frac{1}{1 - \Delta\rho} ; \quad \Delta\rho = 0.0026 \frac{M_t^2}{M_Z^2} - 0.0015 \ln \left(\frac{M_H}{M_W} \right)$

Higgs-search in e^+e^- annihilation: indirect

insertion of running couplings into "Born"-approximation :

partial decay widths of Z: $\Gamma_f = \frac{G_f M_z^3}{6\pi\sqrt{2}} [g_{a,f}^2 + g_{v,f}^2]$ (and thus, also the

cross sections) become dependent on:

- M_t
- M_H
- α_s


\Rightarrow indirect determination (fit) of M_t , M_H , und α_s from combination of all available electro-weak observables
(differential cross sections, partial decay widths, forward-backward asymmetries, τ -polarisation, ...)

$$g_{a,f} = I_{3,f} \quad (3. \text{ Komponente schw. Isospin; } = \pm 1/2)$$

$$g_{v,f} = I_{3,f} - 2Q \sin^2 \theta_w$$

Higgs-search in e^+e^- annihilation: indirect

indirect
(adjusting
radiative corrections):
 $M_H = 84^{+34}_{-26}$ GeV
(68% c.l.)




$114.1 \text{ GeV} < M_H < 154 \text{ GeV}$ (1-sided 95% c.l.)

$M_H < 185 \text{ GeV}$ (incl. 114 GeV lower limit)

n.b.: at the end of LEP (2000), indication for few events with $M_H \sim 115 \text{ GeV}$ (~ 2.3 std. dev.)

Higgs-Search at Hadron colliders: Tevatron




$M_H < \sim 135$ GeV: dominant decay $H \rightarrow b\bar{b}$ ($\sim 90\%$)
 $H \rightarrow \tau^+\tau^-$ ($\sim 8\%$)

$M_H > \sim 135$ GeV: dominant decay $H \rightarrow W^+W^-$

hadron collider: $b\bar{b}$ background from QCD processes
dominates; irreducible;
 $\Rightarrow g g \rightarrow H \rightarrow b\bar{b}$ cannot be used

therefore:


focus on associate production
(ZH , WH) and analyse
e.g. $Z \rightarrow l^+l^-$; $H \rightarrow bb$



$\tau\tau$ decay suitable for all production
channels


Higgs-Search at Hadron colliders: Tevatron

example: $M_H = 120 \text{ GeV}$ and 30 fb^{-1} (model study!)



very difficult measurement; background must be known extremely well!

Collider Run II Integrated Luminosity (delivered)



Tevatron was shut down on Sept. 29, 2011, after 26 years of colliding p and \bar{p}

definition of: significance of signal

● definition of significance

- N_s : number of signal events
 - N_b : number of background events
- $\sqrt{N_b}$ = uncertainty on number of background events

$$S = \frac{N_s}{\sqrt{N_b}}$$


● discovery: $S > 5$

- probability to observe background fluctuation:
- $$\approx 10^{-7}$$

● aim for high significance

- minimize mass resolution σ_M :
- maximize luminosity L :

→ same dependence of efficiency




$$S \sim 1/\sqrt{\sigma_M}$$

$$S \sim \sqrt{L}$$

$m_{\gamma\gamma}$

Expected Tevatron sensitivity

2xCDF Preliminary Projection





Tevatron at end-of-run (Sept. 2011): $< 12 \text{ fb}^{-1}$ / experiment

→ exclusion (3σ) expected: 100 - 117 und 150 - 179 GeV;
 5σ discovery possible to be reached at 160 - 167 GeV.


H \rightarrow WW \rightarrow l ν l ν

- Basic event selection is two reconstructed leptons and missing E_T
- Presence of two neutrinos in final state prevents complete Higgs mass reconstruction
- Separate potential signal from large backgrounds using kinematic event information




$H \rightarrow WW \rightarrow l\nu l\nu$

- Results from thirteen independent search samples are combined to obtain the best possible sensitivity
- No significant, observed excesses in data above predicted SM background contributions




H \rightarrow bb


- Tevatron searches in this decay mode are still the world's most sensitive
- Basic event selection is 0, 1, or 2 leptons and/or missing E_T plus two high E_T jets
- Challenge is separating the small number of potential signal events from the much larger SM background contributions




$ZH \rightarrow wbb$



$WH \rightarrow lbb$



$ZH \rightarrow llbb$



Focus on


Increasing lepton reconstruction and selection efficiencies

Improving the efficiency for tagging bottom quark jets


Optimizing dijet mass resolution

CDF/D0 combined conclusion (July 2012):

- SM Higgs exclusion in the range 147-180 (and 100-103) GeV @95% CL
- Expected exclusion range 139-184 GeV
- 2.5 σ excess in region 115-135 GeV (3.0 σ at $M_H=125$ GeV)



observed
signal strength:



Summary SM Higgs-search w/o LHC data (2012):

Precision measurements of electroweak observables, accounting for radiative corrections ($\propto \log m_H^2$):

$$m_H = 94^{+20}_{-24} \text{ GeV (68% C.L.)}$$

$$m_H < 171 \text{ GeV (95% C.L.)}$$


Direct searches at LEP:

$$m_H > 114.4 \text{ GeV at (95% C.L.)}$$


Direct searches at Tevatron:

$$m_H < 147 \text{ GeV at (95% C.L.)}$$

$$m_H > 180 \text{ GeV at (95% C.L.)}$$



Higgs-search and discovery at LHC (preview):



(see next lecture; 22.1.2018)

Literatur:

- Higgs Particle: The Origin of Mass. [Yasuhiro Okada](#) ([KEK, Tsukuba](#) & [Tsukuba, Graduate U. Adv. Studies](#)) . KEK-TH-1171, Aug 2007. 13pp. e-Print: [arXiv:0708.2016](#) [hep-ph]
- Higgs Boson Properties in the Standard Model and its Supersymmetric Extensions. [John R. Ellis](#) ([CERN](#)) , [Giovanni Ridolfi](#) ([INFN, Genoa](#) & [Genoa U.](#)) , [Fabio Zwirner](#) ([Padua U.](#) & [INFN, Padua](#)) . CERN-PH-TH-2007-012, Feb 2007. Published in **Comptes Rendus Physique** 8:999-1012,2007. e-Print: [hep-ph/0702114](#)
- The Tevatron Higgs exclusion limits and theoretical uncertainties: a critical appraisal. [J. Baglio](#), [A. Djouadi](#), [S. Ferrag](#), [R.M. Godbole](#), . CERN-PH-TH-2010-315, LPT-ORSAY-10-107, Jan 2011. 4pp. [Temporary entry](#) e-Print: [arXiv:1101.1832](#) [hep-ph]
- Updated Combination of CDF and D0 Searches for Standard Model Higgs Boson Production with up to 10.0 fb⁻¹ of Data, <http://arxiv.org/abs/1207.0449>.