

Multivariate analysis of $H \rightarrow b\bar{b}$ in $t\bar{t}H$ associated production mode in ATLAS with fast simulation

Sergey Kotov

MPI für Physik, München

MPI ATLAS meeting, Apr 18, 2005





Max _____
Planck _____
Institut für Physik _____

- 1 Low mass SM Higgs boson overview
- 2 Channel description
- 3 Event reconstruction
- 4 Neural network training
- 5 Analysis results
- 6 Summary

Low mass SM Higgs boson overview

LEP2 experimental bounds on Higgs mass

- precision measurements of EW observables:
 $m_H = 117^{+67}_{-45} \text{ GeV}$
- direct searches: $m_H > 114 \text{ GeV}$



Signature channels for low mass SM Higgs


- $H \rightarrow \tau^+\tau^-$ in VBF production mode
- $H \rightarrow WW^* \rightarrow l\nu l\nu$ in VBF production mode
- $H \rightarrow ZZ^* \rightarrow 4l$ in VBF production mode
- $H \rightarrow \gamma\gamma$ in gluon fusion production mode
- $H \rightarrow b\bar{b}$ in $t\bar{t}H$ associated production mode

Channel description

Features of $t\bar{t}H, H \rightarrow b\bar{b}$ channel

- Complex final state
 - ▶ 6 jets: 4 b-jets and 2 light jets
 - ▶ 1 high- p_t lepton (trigger)
 - ▶ missing energy E_t^{miss} from neutrino
 - ▶ additional jets from ISR/FSR
- Large backgrounds
 - ▶ combinatorial from mis-pairing of b-jets in signal events
 - ▶ irreducible from $t\bar{t}b\bar{b}$ events
 - ▶ reducible from $t\bar{t}jj$ events


full reconstruction of event and very good b-jet tagging are needed to suppress backgrounds




$t\bar{t}H$ signal process diagram

Signal and background cross sections

| Process | $t\bar{t}H$ ($m_H=120$ GeV) | $t\bar{t}b\bar{b}$ | $t\bar{t}jj$ (≥ 6 jets) |
|---------------------|---------------------------------|--------------------|----------------------------------|
| σ, pb | 0.5 | 9.2 | 72 |




$t\bar{t}b\bar{b}$ background



$t\bar{t}jj$ background

Event topology, preselection and data samples



PYTHIA 6.2 and AcerMC 2.3 programs are used for event generation
ATLFAST package is used for fast simulation of ATLAS detector response

Preselection: event topology cuts

- 1 isolated lepton with $p_t > 20(25)$ GeV for $\mu(e)$ and $\eta < 3$
- ≥ 6 jets with $p_t > 20$ GeV and $\eta < 5$
- ≥ 4 of jets identified as b-jets

Data samples

| Process | $t\bar{t}H$ | $t\bar{t}b\bar{b}$ | $t\bar{t}jj$ |
|----------------------------|-------------|--------------------|--------------|
| Generated events, M | 0.6 | 1 | 1 |
| Preselection efficiency, % | 2.1 | 0.8 | 0.02 |

ATLFAST underestimates the b-tagging efficiency ($\epsilon_b=0.42$ compared to $\epsilon_b=0.60-0.65$ in full simulation), so we use an efficiency scale factor in our final calculations

Main reconstruction challenge

How to reconstruct $t\bar{t}$ -pair from all possible $bl\nu - bjj$ combinations properly with correct assignment of b-jets?


Various approaches to $t\bar{t}$ -pair reconstruction

- ATLAS Technical Design Report: select combination with minimal $\Delta^2 = (m_{bl\nu} - m_t)^2 + (m_{bjj} - m_t)^2$
- a recent improved approach which uses likelihood techniques for reconstruction of $t\bar{t}$ -pair
- **this analysis uses neural network technique for $t\bar{t}$ -pair reconstruction**

Reconstruction strategy with Neural Network

- use events which pass preselection criteria (1 lepton, 4 b-jets, 2 light jets)
- determine p_ν from p_l and p_{miss} using m_W constraint (if fails, use approximation $p_\nu^z = p_l^z$)
- select all possible reconstructed combinations of lepton, neutrino, 2 light jets and 2 b-jets for which the reconstructed invariant masses m_{jj} , $m_{bl\nu}$, and m_{bjj} fit inside some mass windows of W boson and t-quark (30 GeV and 70 GeV respectively)
- feed parameters of these combinations through a neural network (which was trained beforehand on a sample of combinations matched and non-matched to MC generator truth table) and select combination with the highest NN output value
- assign the remaining 2 b-jets to the Higgs boson and plot their invariant mass m_{bb}

Multilayer Perceptron



Input layer

Hidden layer

$$h_j = f(\sum x_i w_{ij} + \theta_j)$$

$$f = \frac{1}{1 + e^{-x}}$$

Output layer


$$O = \sum h_j w_j + \theta_o$$

$$E = \frac{1}{N} \sum (O - T)^2$$


- 14 input variables: 5 invariant mass differences, 8 angular separations in ϕ - η plane, 1 sum of transverse momenta
- **TMultiLayerPerceptron** ROOT built-in class is used as neural network (1 hidden layer with 15 nodes)
- 6000 of matched and 21000 of non-matched combinations were used (with proper weights) to train the neural network

Neural network variables


Invariant mass of two light jets




ϕ - η grid between two light jets




Invariant mass of j**j**b system




Angle in ϕ - η grid between **j**-system and **b**-jet




Mass difference of reconstructed t-quarks from nominal



Total transverse momenta of reconstructed t-quarks




Signal/background separation with neural network



Effectiveness of the neural network for background suppression

| Cut | ϵ_{ttH} | ϵ_{cmbn} | ϵ_{ttbb} | ϵ_{ttjj} |
|-----|------------------|-------------------|-------------------|-------------------|
| 0.5 | 0.768 | 0.238 | 0.209 | 0.163 |
| 0.6 | 0.657 | 0.157 | 0.139 | 0.090 |
| 0.7 | 0.510 | 0.093 | 0.080 | 0.046 |
| 0.8 | 0.330 | 0.044 | 0.038 | 0.023 |

Since there're on average about 9 wrongly paired combinations in a $t\bar{t}H$ event, the signal purity is about 35%



Expected number of events for $L = 30 fb^{-1}$

| Scale factor | 4.15 | 3 | 2 | 1 | LH |
|----------------|-------|-------|-------|-------|-------|
| $\epsilon, \%$ | 2.24 | 1.62 | 1.08 | 0.54 | 1.32 |
| N_{ttH} | 101.5 | 73.4 | 49.0 | 24.5 | 41.9 |
| N_{ttbb} | 583.1 | 421.5 | 281.0 | 140.5 | 164.2 |
| N_{ttjj} | 134.6 | 97.3 | 64.9 | 32.4 | 54.6 |
| S/\sqrt{B} | 3.8 | 3.2 | 2.6 | 1.9 | 2.8 |
| Purity | 0.33 | 0.33 | 0.33 | 0.33 | 0.29 |

Purity of reconstructed $t\bar{t}H$ events (fraction of events with all 4 b-jets correctly assigned) is important for finding of the Higgs mass peak

Conclusions

- a neural network approach was tried for reconstruction of $t\bar{t}$ -pair in $t\bar{t}H$ events produced with fast simulation of the ATLAS detector
- the obtained signal significance is still rather low, in large due to imperfections of the fast simulation algorithms

Current status

- 20k of signal $t\bar{t}H$ and 20k of irreducible $t\bar{t}b\bar{b}$ background events were produced at Garching computing farm (up to digitization step, reconstruction is pending)
- a different MC event generator **Alpgen** instead of **Pythia** will probably be used for generation of reducible $t\bar{t}jj$ background

Future plans

- retry the neural network approach on the abovementioned sample of fully simulated events for Rome physics workshop
- more elaborate consideration of the background sources is need to improve signal significance (ATLFAST program needs to be tuned for this using full simulation data, because it is impossible to produce the required amount of background events with full simulation of the ATLAS detector)