

The $Z \rightarrow \tau\tau \rightarrow (\tau_{jet}\nu_\tau)(l\nu_l\nu_\tau)$ analysis with ATLAS detector with early data

Daniele Capriotti

Max Planck Institut fuer Physik
Werner Heisenberg Institut
Muenchen, 80805, Germany
capriott@mppmu.mpg.de

Bonn DPG Meeting, March 15 - 19, 2010



Overview

- Introduction
- Tau reconstruction and identification with the ATLAS detector
- The $Z \rightarrow \tau\tau$ analysis with early data
- Conclusion

Introduction

Introduction



Tau reconstruction is an important (and challenging!) task:

Understanding of the detector:

- Measurement of the τ energy scale
- Measurement of the E_T^{miss} scale

Physics motivations:

- Measurement of W/Z production cross section
- Discovery of the SM and MSSM Higgs boson in $\tau\tau$ final state



Basic Tau Properties

TAU BRANCHING RATIOS

Leptonic Modes (35%)

$e\nu_e\nu_\tau$ 18%

$\mu\nu_\mu\nu_\tau$ 17%

1 Prong Hadronic Modes (47%)

$\pi^-\nu_\tau$ 11%

$\pi^-\pi^0\nu_\tau$ 25%

$\pi^-\pi^0\pi^0\nu_\tau$ 9%

$\pi^-\pi^0\pi^0\pi^0\nu_\tau$ 1%

$K^- + \text{Neutrals}$ 1.5%

3 Prong Hadronic Modes (15%)

$\pi^-\pi^+\pi^-\nu_\tau$ 9%


$\pi^-\pi^+\pi^-\pi^0\nu_\tau$ 4.5%

$K^-\pi^+\pi^-\nu_\tau$ 0.4%

Other Modes ($\approx 3\%$)


TAU CHARACTERISTICS

- $m_\tau \approx 1.8 \text{ GeV}$
- $c\tau = 87 \text{ } \mu\text{m}$
- Hadronic decays are well collimated collection of charged and neutral pions/kaons: leading particle direction reproduces τ direction well
- 1 or 3 charged tracks in the τ -jet



Tau reconstruction and identification with the ATLAS detector

The ATLAS detector at LHC



Tau reconstruction in ATLAS

- ⇒ Leptonic τ decays cannot really be separated from electrons and muons from primary vertex.
- ⇒ The τ reconstruction and ID refer to hadronically decaying τ leptons.


Track seed algorithm

- 1 Seed track with $p_T > 6 \text{ GeV}$ with good quality criteria.
- 2 Association with other tracks in $\Delta R < 0.2$ with quality criteria.
- 3 Total charge $|Q| = 1$.
- 4 Energy flow algorithm

Calorimeter seed algorithm



- 1 Jets with $E_T > 10 \text{ GeV}$.
- 2 Tracks associated if $\Delta R < 0.3$, passing minimal quality criteria.
- 3 Energy calculation summing the weighted calorimeter cells in $\Delta R < 0.4$

Calorimeter + Track seeds matched if $\Delta R < 0.2$



Tau Identification with early data: Safe Identification

- "Safe-variables": discriminating variables for the τ -ID in the early data ("Safe" $\Rightarrow$ small sensitivity to detector-related systematic uncertainties).
- Not used with early data: variables based on: precision tracking, π^0 reconstruction ...

Electromagnetic Radius R_{em} Ratio E_T/p_T of the leading track

$$R_{em} = \frac{\sum_{i=1}^{\Delta R < 0.4} E_{T,i} \sqrt{(\eta_i - \eta)^2 + (\phi_i - \phi)^2}}{\sum_{i=1}^{\Delta R < 0.4} E_{T,i}}$$



i calorimetric cell, E_T η_i ϕ_i measured in the EM calorimeter, η ϕ seeded tau coordinates.

$$E_T^{total}/p_{T1} = \frac{\sum_{i=1} E_{T,i}^{EM} + \sum_{i=1} E_{T,i}^{Had}}{p_{T1}}$$

i calorimetric cell, p_{T1} momentum of the leading track

Taujet efficiency and QCD jet rejection

"Safe Cut" with Calorimeter+Track seed



- Tight, medium, loose cuts corresponding to 0.3, 0.5, 0.7 τ -jet efficiency
- Identification criteria optimized separately for 5 p_T bins (10-25 GeV, 25-45 GeV, 45-70 GeV, 70-100 GeV, > 100 GeV) separately for 1-prong or 3-prong candidates.

The $Z \rightarrow \tau\tau$ selection

$Z \rightarrow \tau\tau$ in early data

Goal: reconstruct a clean sample of $Z \rightarrow \tau\tau$ events. This allows for:

- understanding of the τ -jet reconstruction
- understanding of the background in SM and MSSM Higgs search

Presented analysis performed in order to deal with early data (100 pb^{-1}) at 10 TeV.

	$\sigma \text{ (nb)}$	ϵ_{filter}	Nr. events for 100 pb^{-1}
$Z \rightarrow \tau^+\tau^-$	1.128 (LO)	1	112800
$Z \rightarrow e^+e^-$	1.144 (LO)	0.96	109824
$Z \rightarrow \mu^+\mu^-$	1.144 (LO)	0.96	109824
$W \rightarrow e\nu_e$	11.765 (LO)	0.88	1035320
$W \rightarrow \mu\nu_\mu$	11.765 (LO)	0.88	1035320
$W \rightarrow \tau_{lep}\nu_\mu$	4.148 (LO)	0.87	360876
$W \rightarrow \tau_{had}\nu_\mu$	7.690 (LO)	1	769000
$t\bar{t}$	0.374 (NLO)	0.55	20570
QCD dijet (1e filter) p_T 17-35 GeV	$8.668 \cdot 10^5$	$1.09 \cdot 10^{-3}$	$94.5 \cdot 10^6$
QCD dijet (1e filter) p_T 35-70 GeV	$5.601 \cdot 10^4$	$5.45 \cdot 10^{-3}$	$30.5 \cdot 10^6$
QCD dijet (1 μ filter) p_T 17-35 GeV	$8.668 \cdot 10^5$	$1.02 \cdot 10^{-3}$	$88.4 \cdot 10^6$
QCD dijet (1 μ filter) p_T 35-70 GeV	$5.601 \cdot 10^4$	$5.11 \cdot 10^{-3}$	$28.6 \cdot 10^6$

- samples analysed with full detector simulation
- factorization of QCD background

Selection of the $Z \rightarrow \tau\tau \rightarrow (\tau_{jet}\nu_\tau)(l\nu_l\nu_\tau)$ signature

Medium Electrons

- $p_T > 15 \text{ GeV}$, $|\eta| < 2.5$ and $|Q| = 1$
- Identification flag: medium
- Isolation: E_T (in cone $\Delta R < 0.40$) / $p_T < 0.12$.


Combined muons

- $p_T > 15 \text{ GeV}$, $|\eta| < 2.5$ and $|Q| = 1$
- Identification: combined (inner tracker + muon spectrometer)
- Isolation: E_T (in cone $\Delta R < 0.40$) / $p_T < 0.10$ and no tracks in cone 0.40.


TauJets

- $E_T > 20 \text{ GeV}$, $|\eta| < 2.5$ and $|Q| = 1$
- Tau identification with tight safe variables
- Remove overlap between taujets and combined muons (and medium electron) in $\Delta R < 0.30$
- Electron veto

• Efficiency after selection criteria on the left



• Efficiency after selection criteria on the left



Signal Selection


- ⇒ Combining 1 taujet and 1 lepton (e, μ) with opposite charge ("OppositeSign")
- ⇒ Suppression of $W \rightarrow e(\mu)\nu$ background:

- cut on the transverse invariant mass

$$m_T^{lep, miss} = \sqrt{2p_T^{lep} E_T^{miss} (1 - \cos \phi_{lep, miss})}$$



- cut on the collinearity between lepton and missing energy
- cut on the ΔR between lepton and τ -jet

- ⇒ Counting events in the visible invariant mass windows (M_{vis} , invariant mass of lepton and τ -jet)
- ⇒ Background control with same-sign (τ , lepton) selection ("SameSign").





Results for the $Z \rightarrow \tau\tau \rightarrow (\tau_{jet}\nu_\tau)(e\nu_e\nu_\tau)$ analysis

	$Z \rightarrow \tau^+\tau^-$	$Z \rightarrow e^+e^-$	$W \rightarrow e\nu$	$W \rightarrow \tau_1\nu$	$W \rightarrow \tau_h\nu$	$t\bar{t}$	QCD
Presel $1e+1\tau_{jet}$	450$\pm$7	416 $\pm$ 11	772 $\pm$ 22	52 $\pm$ 7	6 $\pm$ 3	91 $\pm$ 2	235 $\pm$ 46
Opposite Sign							
$1e+1\tau_{jet}$	442$\pm$7	354 $\pm$ 10	580 $\pm$ 19	37 $\pm$ 6	4 $\pm$ 3	79 $\pm$ 2	120 $\pm$ 33
$m_{T^{e,miss}} < 50$ GeV	431$\pm$7	337 $\pm$ 10	141 $\pm$ 10	31 $\pm$ 5	4 $\pm$ 3	24 $\pm$ 1	120 $\pm$ 33
angular correl > -0.15	408$\pm$7	248 $\pm$ 8	50 $\pm$ 6	17 $\pm$ 4	2 $\pm$ 2	19 $\pm$ 1	86 $\pm$ 22
$2.1 < \Delta R < 4.1$	368$\pm$7	232 $\pm$ 8	43 $\pm$ 6	14 $\pm$ 4	2 $\pm$ 2	9 $\pm$ 1	73 $\pm$ 19
$p_T^{ele} < 35$ GeV	330$\pm$6	71 $\pm$ 4	20 $\pm$ 4	9 $\pm$ 3	2 $\pm$ 2	3.0 $\pm$ 0.4	71 $\pm$ 19
$p_T^{\tau_{jet}} < 60$ GeV	326$\pm$6	65 $\pm$ 4	18 $\pm$ 3	8 $\pm$ 3	2 $\pm$ 2	2.7 $\pm$ 0.3	69 $\pm$ 19
$35 < M_{vis} < 80$ GeV	317$\pm$6	11 $\pm$ 2	16 $\pm$ 3	6 $\pm$ 2	2 $\pm$ 2	1.7 $\pm$ 0.3	60 $\pm$ 18
Same Sign							
$1e+1\tau_{jet}$	7$\pm$1	62 $\pm$ 4	192 $\pm$ 11	15 $\pm$ 4	2 $\pm$ 2	12 $\pm$ 1	115 $\pm$ 32
$m_{T^{e,miss}} < 50$ GeV	7$\pm$1	57 $\pm$ 4	35 $\pm$ 5	8 $\pm$ 3	2 $\pm$ 2	4.0 $\pm$ 0.4	114 $\pm$ 32
angular correl > -0.15	5$\pm$1	41 $\pm$ 3	18 $\pm$ 3	3 $\pm$ 2	<1.6	3.3 $\pm$ 0.4	84 $\pm$ 23
$2.1 < \Delta R < 4.1$	4$\pm$1	30 $\pm$ 3	12 $\pm$ 3	2 $\pm$ 1	<1.6	1.7 $\pm$ 0.4	66 $\pm$ 17
$p_T^{ele} < 35$ GeV	3$\pm$1	8 $\pm$ 1	7 $\pm$ 2	2 $\pm$ 1	<1.6	0.3 $\pm$ 0.1	64 $\pm$ 17
$p_T^{\tau_{jet}} < 60$ GeV	3$\pm$1	8 $\pm$ 1	7 $\pm$ 2	2 $\pm$ 1	<1.6	0.3 $\pm$ 0.1	62 $\pm$ 17
$35 < M_{vis} < 80$ GeV	2$\pm$1	4 $\pm$ 1	5 $\pm$ 2	2 $\pm$ 1	<1.6	0.2 $\pm$ 0.1	55 $\pm$ 16



Results for the $Z \rightarrow \tau\tau \rightarrow (\tau_{jet}\nu_\tau)(\mu\nu_\mu\nu_\tau)$ analysis

	$Z \rightarrow \tau^+\tau^-$	$Z \rightarrow \mu^+\mu^-$	$W \rightarrow \mu\nu$	$W \rightarrow \tau_\mu\nu$	$W \rightarrow \tau_h\nu$	$t\bar{t}$	QCD
Presel $1\mu+1\tau_{jet}$	600±8	143±6	906±24	62±8	<1.6	97±2	179±37
OppositeSign							
$1\mu+1\tau_{jet}$	590±8	102±5	689±21	47±7	<1.6	84±2	95±28
$m_T^{\mu,miss} < 50 \text{ GeV}$	573±8	74±4	157±10	36±6	<1.6	24±1	95±28
angular correl > -0.15	537±8	49±3	55±6	18±4	<1.6	19±1	67±19
$2.1 < \Delta R < 4.1$	489±8	43±3	46±6	17±4	<1.6	9±1	55±16
$p_T^{ele} < 35 \text{ GeV}$	444±7	14±2	26±4	15±4	<1.6	3.4±0.4	54±16
$p_T^{\tau_{jet}} < 60 \text{ GeV}$	440±7	13±2	25±4	13±3	<1.6	2.8±0.3	53±16
$35 < M_{vis} < 80 \text{ GeV}$	430±7	10±1	22±4	12±3	<1.6	2.0±0.3	48±15
SameSign							
$1\mu+1\tau_{jet}$	10±1	41±3	216±12	15±4	<1.6	12±1	84±24
$m_T^{\mu,miss} < 50 \text{ GeV}$	10±1	27±2	49±6	9±3	<1.6	4.6±0.4	84±24
angular correl > -0.15	7±1	17±2	22±4	5±2	<1.6	3.4±0.4	59±17
$2.1 < \Delta R < 4.1$	6±1	13±2	18±4	3±2	<1.6	1.8±0.4	48±14
$p_T^{ele} < 35 \text{ GeV}$	5±1	4±1	7±2	3±2	<1.6	0.4±0.1	48±14
$p_T^{\tau_{jet}} < 60 \text{ GeV}$	4±1	4±1	6±2	3±2	<1.6	0.3±0.1	47±13
$35 < M_{vis} < 80 \text{ GeV}$	4±1	3±1	5±2	3±2	<1.6	0.3±0.1	42±13




Conclusion

- $Z \rightarrow \tau\tau$ analysis optimized for early data (100 pb^{-1}): good identification of the τ -jets and separation from the QCD background
- Data driven estimation of the QCD background from the Same Sign control data sample
- Extend the analysis to the $Z + \text{jet}$ final state (dominant background for SM Higgs searches in the $\tau\tau$ final state)


Backup Slides

- Safe variables for tau identification
- Factorization of the QCD background
- Transverse mass with missing energy and angular correlation
- Results with loose lepton isolation

Safe variables for 1 prong taujets



Safe variables for 3 prong taujets



Factorization of the QCD background

Due to low MC statistics for QCD background, its contribution has to be estimated from factorization procedure.

Procedure:



- ➊ Performing the analysis with no safe ID cuts on τ -jets and avoiding lepton isolation (cut on E_T/p_T and number of tracks in cone)
- ➋ Efficiencies of the lepton isolation and τ -jet identification are calculated separately: no correlation found with the other cuts applied in the analysis
- ➌ QCD events weighted by the efficiency of the lepton isolation and τ -jet identification

	QCD pT: 17-35 GeV, 1 electron filter	QCD pT: 35-70 GeV, 1 electron filter
lepton isolation efficiency	0.0359 ± 0.0110	0.0224 ± 0.0032
τ -jet identification efficiency	0.0229 ± 0.0087	0.0149 ± 0.0026
	QCD pT: 17-35 GeV, 1 muon filter	QCD pT: 35-70 GeV, 1 muon filter
lepton isolation efficiency	0.0158 ± 0.0048	0.0024 ± 0.0005
τ -jet identification efficiency	0.0274 ± 0.0063	0.0140 ± 0.0012

Transverse invariant mass

- Transverse invariant mass lepton + missing energy:

$$m_T = \sqrt{2p_T^{lep} E_T^{miss} (1 - \cos \phi_{lep,miss})}$$




- $m_T^{lep,miss} < 50 \text{ GeV}$ for the analysis


Angular Correlation

- Angular Correlation: $\Delta\psi = \cos(\phi_{lep} - \phi_{miss}) + \cos(\phi_{jet} - \phi_{miss})$

Electron+TauJet





Muon+TauJet



- $\Delta\psi > -0.15$

Distance between lepton- τ_{jet}

- Distance between the two visible particles: $\Delta R = \sqrt{\Delta\phi^2 + \Delta\theta^2}$



- $2.14 < \Delta R_{lep, \tau_{jet}} < 4.14$ for the analysis


Old results for the $Z \rightarrow \tau\tau \rightarrow e\tau_{jet}$ analysis

Looser lepton isolation in order to compare SS and OS events with higher statistics


Number of events shown after trigger selection [trigger efficiency in brackets]

	$Z \rightarrow \tau^+\tau^-$	$Z \rightarrow e^+e^-$	$W \rightarrow e\nu$	$W \rightarrow \tau\nu$	$W \rightarrow \tau_h\nu$	$t\bar{t}$	QCD
Presel $1e+1\tau_{jet}$	508$\pm$8	746 $\pm$ 14	764 $\pm$ 22	60 $\pm$ 7	6 $\pm$ 3	80 $\pm$ 2	1759 $\pm$ 256
Opposite Sign (OS)							
$1e+1\tau_{jet}$	497$\pm$7	668 $\pm$ 14	567 $\pm$ 19	47 $\pm$ 7	4 $\pm$ 3	69 $\pm$ 2	908 $\pm$ 134
$m_T^{e,miss} < 35$ GeV	457$\pm$7	511 $\pm$ 12	78 $\pm$ 7	31 $\pm$ 5	4 $\pm$ 3	14 $\pm$ 1	885 $\pm$ 130
$2.1 < \Delta R < 4.1$	415$\pm$7	464 $\pm$ 11	61 $\pm$ 6	23 $\pm$ 5	4 $\pm$ 3	6.6 $\pm$ 0.5	714 $\pm$ 105
$p_T^{ele} < 35$ GeV	385$\pm$7	182 $\pm$ 7	49 $\pm$ 6	19 $\pm$ 4	4 $\pm$ 3	2.9 $\pm$ 0.3	700 $\pm$ 103
$p_T^{\tau_{jet}} < 60$ GeV	380$\pm$6	161 $\pm$ 7	45 $\pm$ 5	17 $\pm$ 4	4 $\pm$ 3	2.4 $\pm$ 0.3	667 $\pm$ 97
$35 < M_{vis} < 80$ GeV	369$\pm$6 [0.97]	24 $\pm$ 3 [1]	38 $\pm$ 5 [1]	15 $\pm$ 4 [1]	4 $\pm$ 3 [1]	1.5 $\pm$ 0.3 [1]	550 $\pm$ 80 [0.99]
Same Sign (SS)							
$1e+1\tau_{jet}$	10$\pm$1	78 $\pm$ 5	197 $\pm$ 11	13 $\pm$ 3	2 $\pm$ 2	11 $\pm$ 1	850 $\pm$ 126
$m_T^{e,miss} < 35$ GeV	9$\pm$1	61 $\pm$ 4 [	18 $\pm$ 3	6 $\pm$ 2	<1.6	2.8 $\pm$ 0.3	831 $\pm$ 123
$2.1 < \Delta R < 4.1$	7$\pm$1	44 $\pm$ 3	13 $\pm$ 3	3 $\pm$ 2	<1.6	1.3 $\pm$ 0.2	654 $\pm$ 98
$p_T^{ele} < 35$ GeV	7$\pm$1	16 $\pm$ 2	10 $\pm$ 3	3 $\pm$ 2	<1.6	0.4 $\pm$ 0.1	639 $\pm$ 96
$p_T^{\tau_{jet}} < 60$ GeV	6$\pm$1	15 $\pm$ 2	10 $\pm$ 3	3 $\pm$ 2	<1.6	0.4 $\pm$ 0.1	610 $\pm$ 91
$35 < M_{vis} < 80$ GeV	6$\pm$1 [1]	7 $\pm$ 1 [1]	7 $\pm$ 2 [1]	3 $\pm$ 2 [1]	<1.6	0.3 $\pm$ 0.1 [1]	505 $\pm$ 75 [0.99]

Electron+TauJet Opposite Sign



Electron+TauJet Same Sign




Old results for the $Z \rightarrow \tau\tau \rightarrow \mu\tau_{jet}$ analysis

Looser lepton isolation in order to compare SS and OS events with higher statistics

Number of events shown after trigger selection [trigger efficiency in brackets]

	$Z \rightarrow \tau^+\tau^-$	$Z \rightarrow \mu^+\mu^-$	$W \rightarrow \mu\nu$	$W \rightarrow \tau_l\nu$	$W \rightarrow \tau_h\nu$	$t\bar{t}$	QCD
Presel $1\mu+1\tau_{jet}$	613$\pm$8	134 $\pm$ 5	901 $\pm$ 24	57 $\pm$ 7	<1.6	106 $\pm$ 2	2685 $\pm$ 407
Opposite Sign (OS)							
$1\mu+1\tau_{jet}$	602$\pm$8	96 $\pm$ 5	674 $\pm$ 21	44 $\pm$ 6	<1.6	91 $\pm$ 2	1373 $\pm$ 208
$m_T^{\mu,miss} < 35$ GeV	540$\pm$8	52 $\pm$ 3	66 $\pm$ 7	29 $\pm$ 5	<1.6	17 $\pm$ 1	1317 $\pm$ 199
$2.1 < \Delta R < 4.1$	492$\pm$7	47 $\pm$ 3	53 $\pm$ 6	24 $\pm$ 5	<1.6	9 $\pm$ 1	1062 $\pm$ 162
$p_T^{ele} < 35$ GeV	457$\pm$7	18 $\pm$ 2	40 $\pm$ 5	21 $\pm$ 4	<1.6	4.2 $\pm$ 0.4	1049 $\pm$ 159
$p_T^{\tau_{jet}} < 60$ GeV	452$\pm$7	17 $\pm$ 2	37 $\pm$ 5	20 $\pm$ 4	<1.6	3.3 $\pm$ 0.4	972 $\pm$ 146
35 < $M_{vis} < 80$ GeV	440$\pm$7 [0.86]	13 $\pm$ 2 [0.93]	30 $\pm$ 4 [0.97]	18 $\pm$ 4 [0.95]	<1.6	2.2 $\pm$ 0.3 [1]	771 $\pm$ 114 [0.80]
Same Sign (SS)							
$1\mu+1\tau_{jet}$	11$\pm$1	38 $\pm$ 3	226 $\pm$ 12	13 $\pm$ 3	<1.6	14 $\pm$ 1	1312 $\pm$ 201
$m_T^{\mu,miss} < 35$ GeV	9$\pm$1	16 $\pm$ 2	22 $\pm$ 4	4 $\pm$ 2	<1.6	2.6 $\pm$ 0.3	1254 $\pm$ 191
$2.1 < \Delta R < 4.1$	7$\pm$1	10 $\pm$ 1	17 $\pm$ 3	2 $\pm$ 1	<1.6	1.3 $\pm$ 0.2	1014 $\pm$ 156
$p_T^{ele} < 35$ GeV	6$\pm$1	5 $\pm$ 1	12 $\pm$ 3	2 $\pm$ 1	<1.6	0.5 $\pm$ 0.1	1000 $\pm$ 153
$p_T^{\tau_{jet}} < 60$ GeV	5$\pm$1	5 $\pm$ 1	11 $\pm$ 3	2 $\pm$ 1	<1.6	0.4 $\pm$ 0.1	921 $\pm$ 140
35 < $M_{vis} < 80$ GeV	5$\pm$1	4 $\pm$ 1 [1]	8 $\pm$ 2 [1]	2 $\pm$ 1 [0.67]	<1.6	0.3 $\pm$ 0.1 [1]	745 $\pm$ 112 [0.80]

Muon+TauJet Opposite Sign



Muon+TauJet Same Sign

