

# Background studies for $B^0$ mixing with hadronic final states at the Belle II experiment

Caspar Schmitt  
cschmitt@mpp.mpg.de


DPG-Verhandlung 2021, Dortmund



# Cabibbo-Kobayashi-Maskawa Matrix

Unitary CKM-matrix relates weak quark eigenstates to strong quark eigenstates and governs flavor-changing quark-transitions.

Unitarity yields 6 relations, i.e. unitarity triangles, that serve as SM precision test. Non-trivial angles indicate CP-violation.



CKM fitter

# Motivation

All-hadronic charmed  $B \rightarrow \pi D$  decays have high branching ratios and provide high statistics.

This talk focusses on estimating **systematic uncertainties** related to backgrounds in measurements of the **mixing frequency**  $\Delta m_d$  and the **lifetime**  $\tau_{B^0}$ . These time-dependent measurements are **systematically limited** at B-factories.

Given the large statistics, we expect to be able to make precise measurements with only  $64\text{fb}^{-1}$  of data available<sup>1</sup>.


---

<sup>1</sup>Belle II note on time-dependent CP-violation and mixing

# Time-dependent measurements at B-factories

Electron-positron collider KEK at  $Y(4S)$  resonance produces coherent B meson pairs at record luminosities with low backgrounds.


Proper-time difference measurement of B meson pairs yields lifetime. Here, we neglect detector resolution, wrong tags and CP-violation in mixing.



$$N(\Delta t) = \frac{1}{2\tau_{B^0}} \cdot \exp\left[\frac{-|\Delta t|}{\tau_{B^0}}\right]$$

# Mixing measurements at B-factories

The B mixing parameter  $\Delta m_d$  can be extracted from the mixing asymmetry in flavor-specific decays<sup>1</sup>.





$$A_{mix}(\Delta t) = \frac{P_{OF}(\Delta t) - P_{SF}(\Delta t)}{P_{OF}(\Delta t) + P_{SF}(\Delta t)} = \cos(\Delta m_d \Delta t)$$

$$\text{with } P_{OF(SF)} = \exp\left[\frac{-|\Delta t|}{4\tau_{B^0}}\right] \cdot [1 + (-) \cos(\Delta m_d \Delta t)]$$

Uncertainties on  $\Delta t$ ,  $\Delta m_d$  and peaking backgrounds enter as systematic uncertainties.

# MC studies of backgrounds in $B \rightarrow \pi D(\rightarrow K \pi \pi)$

Signal B decays peak in both the energy difference  $\Delta E \equiv E_B^{cms} - E_{beam}^{cms}$  and in the beam-energy constrained mass  $M_{bc} \equiv \sqrt{(E_{beam}^{cms})^2 - (p_B^{cms})^2}$ .



continuum backgrounds

combinatorial crossfeed

combinatorial self-crossfeed with signal decay

$q\bar{q}$

peaking backgrounds



$B \rightarrow DK$

partly reconstructed  $B \rightarrow \pi D^*$ ,  $B \rightarrow \rho D$

mis-identified  $e, \mu$  as  $\pi$

# Fit on $64\text{fb}^{-1}$ Belle II data

We fit the **fixed shapes from MC** on data in **two slices in  $M_{bc}$** . We fix the relative yields of crossfeed to self-crossfeed and of  $B \rightarrow DK$  to other peaking backgrounds.




yields normed to total	data	MC truth
crossfeed	$1840 \pm 250$	$1747 \pm 11$
$q\bar{q}$	$5460 \pm 110$	$5485 \pm 19$
$B \rightarrow KD$	$8 \pm 9$	$204 \pm 4$
signal	$7150 \pm 110$	$6732 \pm 21$

## Why no 2D-fit?

To further increase the distinction power of the fit, a full 2D-fit in the  $(\Delta E, M_{bc})$  plane would be useful.


However, we find **non-negligible correlations** between the two variables, which forbids an independent modelling of the shapes. For peaking components, the Pearson correlation coefficients are  $\mathcal{O}(0.2) \gg 0$ .






## MC studies of backgrounds in $\Delta t$

We aim to constrain the continuum backgrounds in  $\Delta t$  from the sidebands  $M_{bc} < 5.265\text{GeV}$  and  $|\Delta E| > 0.05\text{GeV}$ . In both the crossfeed and  $q\bar{q}$  background components, we find that the shapes in sideband and signal region allow for interpolation.



crossfeed in signal region and sideband.



$q\bar{q}$  in signal region and sideband.

## Summary and outlook

- ▶ Study of systematic uncertainties related to backgrounds in time-dependent measurements of mixing frequency  $\Delta m_d$  and lifetime  $\Delta t_{B^0}$  at B-factories.
- ▶ MC studies of background shapes in  $B \rightarrow \pi D(\rightarrow K \pi \pi)$  decay mode.
- ▶ Fit shapes from MC fitted onto data in two slices in  $M_{bc}$ . Good discrimination in continuum backgrounds, while non-negligible correlations in peaking backgrounds.
- ▶ MC background shapes in  $\Delta t$  look promising for background constraint from sideband.

### Outlook

- ▶ Extend analysis to multiple decay modes, notably to  $B \rightarrow \pi D^*(\rightarrow \pi_{\text{slow}} D)$ .
- ▶ Analyse data in segments of flavor-tag figure of merit.
- ▶ Evaluate systematic uncertainties related to backgrounds for first full Belle II mixing parameter measurement.